For Research Use Only

GAA Polyclonal antibody

Catalog Number: 14367-1-AP

6 Publications



Basic Information

Catalog Number: GenBank Accession Number: 14367-1-AP BC040431

Size: GeneID (NCBI): 2548

Source: UNIPROT ID:

Rabbit P10253
Isotype: Full Name:

IgG glucosidase, alpha; acid

Immunogen Catalog Number:Calculated MW:AG5690105 kDa

Observed MW: 70-76 kDa

Purification Method: Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

Applications

Tested Applications: IHC, IP, WB, ELISA Cited Applications: WB, IF, IHC Species Specificity: human, mouse

Cited Species:

human, rat, mouse, zebrafish

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: DU 145 cells, mouse liver tissue

IP: mouse liver tissue,

IHC: human pancreas cancer tissue, mouse skeletal

muscle tissue, mouse liver tissue

Background Information

GAA(Alpha-1,4-glucosidase) is a lysosomal enzyme involved in the degradation of glycogen within cellular vacuoles. After translation, GAA undergoes proteolytic processing to form two lengths of lysosomal a-glucosidase, and both N-terminal and C-terminal processing occur. The full sequence can be further processed into a mature form. The 105 kDa GAA can be cleaved into 70 kDa or 76 kDa form.

Notable Publications

Author	Pubmed ID	Journal	Application
Atsushi Kato	35838176	Org Biomol Chem	IF
Atsushi Kato	35072486	J Med Chem	IF
Jing Wu	28045567	DNA Cell Biol	WB

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 14367-1-AP (GAA antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-GAA (IP:14367-1-AP, 4ug; Detection:14367-1-AP 1:600) with mouse liver tissue lysate 4000ug.



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 14367-1-AP (GAA antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).