For Research Use Only

NOP58 Polyclonal antibody

Catalog Number: 14409-1-AP

Featured Product

6 Publications



Basic Information

Catalog Number: GenBank Accession Number: 14409-1-AP BC032592 GeneID (NCBI): Size: 1200 ug/ml 51602 **UNIPROT ID:** Source: Rabbit Q9Y2X3

Isotype: NOP58 ribonucleoprotein homolog

(veast) Immunogen Catalog Number:

AG5740 Calculated MW:

60 kDa Observed MW: 60 kDa

Full Name:

Antigen affinity purification Recommended Dilutions:

WB 1:2000-1:12000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:2000-1:8000

Purification Method:

Applications

Tested Applications: WB, IHC, IP, ELISA **Cited Applications:** WB, IF, CoIP, RIP Species Specificity:

human, mouse, rat **Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, Jurkat cells, K-562 cells, MCF-7 cells

IP: K-562 cells,

IHC: mouse kidney tissue,

Background Information

The box C/D snoRNPs, is named for the short, conserved box C (UGAUGA) and box D (CUGA) sequences present in their RNA moiety. The box C/D sequences are also required for multiple aspects. of snoRNA biogenesis, including: RNA stability, intronic processing, nucleolar targeting, nuclear retention, and 59 trimethylguanosine cap formation [PMID:9649444,8878486]. Nop58 is a component common to the box C/D small nucleolar ribonucleoproteins [PMID:10606270].

Notable Publications

Author	Pubmed ID	Journal	Application
Yick W Fong	25407680	Elife	WB
Beatriz Alvarado-Hernandez	34787459	J Virol	WB,CoIP
Marlene Cervantes	32450515	iScience	CoIP,RIP

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 14409-1-AP (NOP58 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



WB result of NOP58 antibody (14409-1-AP; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-NOP58 transfected HeLa cells.



IP result of anti-NOP58 (IP:14409-1-AP, 4ug; Detection:14409-1-AP 1:500) with K-562 cells lysate 3600ug.



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 14409-1-AP (NOP58 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).