

For Research Use Only

# TCF12/HEB Polyclonal antibody

Catalog Number: 14419-1-AP

Featured Product

12 Publications



## Basic Information

Catalog Number:

14419-1-AP

Size:

400 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG5765

GenBank Accession Number:

BC050556

GeneID (NCBI):

6938

UNIPROT ID:

Q99081

Full Name:

transcription factor 12

Calculated MW:

73 kDa

Observed MW:

73 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:10000

IHC 1:50-1:500

## Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

Positive Controls:

WB : HepG2 cells, Raji cells

IHC : human cervical cancer tissue,

## Background Information

The product of TCF12 gene (TCF12, also called HTF4 or HEB) is a member of helix-loop-helix (HLH) protein family [PMID:22130667]. TCF12 is a transcriptional regulator that involved in the initiation of neuronal differentiation. It's also known as E proteins, and activates transcription by binding to the E box (5'-CANNTG-3'), because of their direct DNA (E-box) binding ability [PMID: 19204326]. TCF12 was shown to suppress E-cadherin expression during the early stages of renal tubular epithelial cell dedifferentiation [PMID:17202424].

## Notable Publications

Author	Pubmed ID	Journal	Application
Sisu Han	30201687	Development	IF
Thorsen Kasper K	21999571	BMC Genomics	IHC
Liang Liu	36377508	CNS Neurosci Ther	WB

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

Selected Validation Data



HepG2 cells were subjected to SDS PAGE followed by western blot with 14419-1-AP (TCF12/HEB antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 14419-1-AP (TCF12/HEB antibody) at dilution of 1:100 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).