For Research Use Only

AMH Polyclonal antibody Catalog Number:14461-1-AP Featured Product

Featured Product






Basic Information	Catalog Number: 14461-1-AP	GenBank Accession N BC049194	lumber:	Purification Method: Antigen affinity purification	
	Concentration:	GenelD (NCBI):		Recommended Dilutions:	
	700 ug/ml	268		WB 1:500-1:1000	
	Source:	UNIPROT ID:		IHC 1:50-1:500 IF-P 1:50-1:500	
	Rabbit	P03971		II-F 1.30-1.300	
	Isotype: IgG	Full Name: anti-Mullerian hormo	one		
	Immunogen Catalog Number: AG5670	Calculated MW: 59 kDa			
		Observed MW: 60-65 kDa			
Applications	Tested Applications:		Positive Con	rrols:	
	WB, IHC, IF-P, ELISA Cited Applications:		WB : human t	estis tissue,	
	WB, IHC, IF		IHC : mouse ovary tissue, human pancreas tissue, rat ovary tissue, human ovary tumor tissue, human testis		
	Species Specificity:		tissue	numan ovary tumor tissue, numan testi	
	human, mouse, rat		IF-P: mouse	ovary tissue,	
	Cited Species:	roat		-	
		human, mouse, rat, pig, chicken, goat Note-IHC: suggested antigen retrieval with			
	TE buffer pH 9.0; (*) Altern retrieval may be performe buffer pH 6.0	natively, antigen			
Background Information	Anti-Müllerian hormone (AMH), also called Müllerian-inhibiting substance (MIS), is a hormone that is best known for its production by fetal testes in mammals and as the inhibitor of Müllerian duct development in males. AMH is also expressed in granulosa cells of preantral and small antral follicles in the ovary. More recently, AMH has been studied for its role in ovarian folliculogenesis and as a potential marker of ovarian reserve. The deduced protein sequence of the AMH monomer contains a 25 amino acid secretion specific signal peptide and a monomeric protein of 535 amino acids that, when glycosylated at two glycosylation sites, has a molecular weight of 65-70 kDa. This antibody is specific to AMH.				
	studied for its role in ovarian foll sequence of the AMH monomer c of 535 amino acids that, when gly	liculogenesis and as a pote ontains a 25 amino acid se	cretion specific	ovarian reserve. The deduced protein signal peptide and a monomeric protei	
	studied for its role in ovarian foll sequence of the AMH monomer c of 535 amino acids that, when gly antibody is specific to AMH.	liculogenesis and as a pote ontains a 25 amino acid se	cretion specific ation sites, has	ovarian reserve. The deduced protein signal peptide and a monomeric protei	
	studied for its role in ovarian foll sequence of the AMH monomer c of 535 amino acids that, when gly antibody is specific to AMH.	iculogenesis and as a pote ontains a 25 amino acid se ycosylated at two glycosyl Pubmed ID Journ	cretion specific ation sites, has	ovarian reserve. The deduced protein signal peptide and a monomeric protei a molecular weight of 65-70 kDa. This	
Notable Publications	studied for its role in ovarian foll sequence of the AMH monomer c of 535 amino acids that, when gly antibody is specific to AMH. Author Tong Wu	iculogenesis and as a pote ontains a 25 amino acid se ycosylated at two glycosyl Pubmed ID Journ 36278634 J Func	ecretion specific ation sites, has al	ovarian reserve. The deduced protein signal peptide and a monomeric protei a molecular weight of 65-70 kDa. This Application	
	studied for its role in ovarian foll sequence of the AMH monomer c of 535 amino acids that, when gly antibody is specific to AMH. Author Tong Wu Lei Wang	Pubmed ID Journ 36525856 Theric	ecretion specific ation sites, has al al	ovarian reserve. The deduced protein signal peptide and a monomeric protei a molecular weight of 65-70 kDa. This Application IHC	


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human testis tissue were subjected to SDS PAGE followed by western blot with 14461-1-AP (AMH antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse ovary tissue slide using 14461-1-AP (AMH antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse ovary tissue slide using 14461-1-AP (AMH antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse ovary tissue using AMH antibody (14461-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).