For Research Use Only

TFE3 Polyclonal antibody

Catalog Number: 14480-1-AP

Featured Product

21 Publications



Basic Information

Catalog Number: GenBank Accession Number: 14480-1-AP BC026027 GeneID (NCBI): Concentration: $600 \mu g/ml$ 7030 **UNIPROT ID:** Source: Rabbit P19532 Full Name: Isotype:

transcription factor binding to IGHM

Immunogen Catalog Number: AG5863 Calculated MW: 62 kDa Observed MW:

70 kDa

enhancer 3

Applications

Tested Applications: WB, IP, IHC, ELISA Cited Applications: WB, IF, IP, ChIP Species Specificity: human, mouse, rat

human, mouse, rat Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Cited Species:

Recommended Dilutions: WB 1:500-1:2000

Purification Method:

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

Positive Controls:

WB: K-562 cells, human brain tissue, mouse spleen

tissue, rat brain tissue IP: K-562 cells.

IHC: human renal cell carcinoma tissue,

Background Information

The protein encoded by this gene is a subunit of the CRSP (cofactor required for SP1 activation) complex, which, along with TFIID, is required for efficient activation by SP1. This protein is also a component of other multisubunit complexes e.g. thyroid hormone receptor-(TR-) associated proteins which interact with TR and facilitate TR function on DNA templates in conjunction with initiation factors and cofactors.

Notable Publications

| Author | Pubmed ID | Journal | Application |
|------------------|-----------|-------------------------------|-------------|
| Bo Wang | 33019842 | Autophagy | WB,IF,IP |
| Sebastian Boland | 36207292 | Nat Commun | WB |
| Xiu-Lei Zhang | 31103748 | Biochim Biophys Acta Gen Subj | WB |

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



K-562 cells were subjected to SDS PAGE followed by western blot with 14480-1-AP (TFE3 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



IP result of anti-TFE3 (IP:14480-1-AP, 4ug; Detection:14480-1-AP 1:500) with K-562 cells lysate 1200ug.



Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 14480-1-AP (TFE3 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).