For Research Use Only

CD34 Polyclonal antibody

Catalog Number: 14486-1-AP 87 Publications



Basic Information

Catalog Number: 14486-1-AP

Concentration: GeneID (NCBI):

550 ug/ml 94

Source: UNIPROT ID:
Rabbit P28906

Isotype: Full Name:
IgG CD34 molecule
Immunogen Catalog Number: Calculated MW:
AG5887 41 kDa

Observed MW: 100-120 kDa

BC039146

GenBank Accession Number:

Purification Method:

Antigen affinity purification Recommended Dilutions:

WB 1:500-1:3000 IHC 1:1000-1:4000 IF-P 1:50-1:500

Applications

Tested Applications: WB, IHC, IF-P, ELISA Cited Applications: WB, IHC, IF Species Specificity:

human
Cited Species:

human, pig, rabbit, bovine, sheep

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HL-60 cells, TF-1 cells

IHC: human tonsillitis tissue, human gliomas tissue, human hysteromyoma tissue, human liver cancer tissue, human placenta tissue

IF-P: human tonsillitis tissue, human liver cancer tissue, human placenta tissue

Background Information

CD34 is a 105- to 120-kDa glycophosphoprotein expressed on the majority of hematopoietic stem/progenitor cells, bone marrow stromal cells, capillary endothelial cells, embryonic fibroblasts, and some nerve tissue. CD34 is a commonly used marker for identifying human hematopoietic stem/progenitor cells and mediates cell adhesion and lymphocyte homing by binding L-selectin and E-selectin ligands. CD34 is also one of the best negative selection markers for characterizing and/or isolating human MSCs from bone marrow and other sources. Along with other positive selection markers (such as CD29, CD44, CD90, CD105 and CD166), negative selection markers (such as CD34 and CD45) are used for MSC identification. The calculated molecular mass of human CD34 is 41 kDa, various forms with different molecular weights may be produced due to different glycosylation patterns and alternative splicing (PMID: 24375067; 15750786).

Notable Publications

Author	Pubmed ID	Journal	Application
Kexin Ma	32961493	Ecotoxicol Environ Saf	WB
Ruidong Wan	34521949	Sci Rep	IHC
Zuping Wu	36068594	Stem Cell Res Ther	FC

Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 14486-1-AP (CD34 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 14486-1-AP (CD34 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 14486-1-AP (CD34 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using 14486-1-AP (CD34 antibody), at dilution of 1:100 and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using 14486-1-AP (CD34 antibody), at dilution of 1:100 and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 14486-1-AP (CD34 antibody) at dilution of 1:6000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).