For Research Use Only

SHARPIN Polyclonal antibody

Catalog Number: 14626-1-AP

Featured Product

107 Publications



Basic Information

Catalog Number:

14626-1-AP

BC025244

Concentration:

650 ug/ml

S1858

Source:

UNIPROT ID:

Rabbit

Q9H0F6

Isotype:

GenBank Accession Number:

BC025244

GeneID (NCBI):

81858

UNIPROT ID:

Q9H0F6

Full Name:

gG SHANK-associated RH domain interactor

Immunogen Catalog Number: Interactor
AG6201 Calculated MW:

40 kDa Observed MW:

40 kDa

Purification Method:

Antigen affinity purification Recommended Dilutions:

WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:100-1:400 IF/ICC 1:200-1:800

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB, IHC, IF, IP, CoIP Species Specificity: human, mouse Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

human, mouse, rat

Positive Controls:

WB: A549 cells, mouse brain tissue, mouse skeletal muscle tissue, Jurkat cells, Raji cells

IP: Raji cells,

IHC: human lymphoma tissue, human small intestine

tissue

IF/ICC: Jurkat cells,

Background Information

SHARPIN is as a third component of the linear ubiquitin chain assembly complex (LUBAC), recruited to the CD40 and TNF receptor signaling complexes together with its other constituents, HOIL1 and HOIP(PMID:21455173). SHARPIN is enriched in the postsynaptic density and forms a complex with SHANK in heterologous cells and brain(PMID:11178875). It has 2 isoforms produced by alternative splicing.

Notable Publications

Author	Pubmed ID	Journal	Application
Matthias Kist	32999468	Cell Death Differ	WB,IP
Lucia Taraborrelli	30254289	Nat Commun	WB
Umar Butt	36148554	J Cell Sci	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 14626-1-AP (SHARPIN antibody) at dilution of 1:40000 incubated at room temperature for 1.5 hours.




IP result of anti-SHARPIN (IP:14626-1-AP, 3ug; Detection:14626-1-AP 1:1500) with Raji cells lysate 2520ug.



Immunohistochemical analysis of paraffinembedded human lymphoma tissue slide using 14626-1-AP (SHARPIN Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lymphoma tissue slide using 14626-1-AP (SHARPIN Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed Jurkat cells using SHARPIN antibody (14626-1-AP) at dilution of 1:400 and Multi-rAb Coralite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).