For Research Use Only

CNPY2/MSAP Polyclonal antibody

Catalog Number: 14635-1-AP

Featured Product

16 Publications



Basic Information

Catalog Number:

GenBank Accession Number:

14635-1-AP

BC065015

Concentration:

300 ug/ml

10330

Source:

UNIPROT ID:

Rabbit

Q9Y2B0

Isotype:

GenBank Accession Number:

UNCBI):

GeneID (NCBI):

Q9Y2B0

Full Name:

gG canopy 2 homolog (zebrafish)

Immunogen Catalog Number:Calculated MW:AG621421 kDa

Observed MW: 17-21 kDa Purification Method: Antigen affinity purification Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:200-1:800 IF/ICC 1:20-1:200

WB 1:1000-1:8000

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), IP, ELISA

Cited Applications: WB, IHC, IF

Species Specificity: human, mouse, rat Cited Species: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:**

WB: HepG2 cells, mouse pancreas tissue, human liver tissue, mouse liver tissue, rat liver tissue

IP: HepG2 cells,

IHC: human liver tissue, human pancreas cancer tissue

IF/ICC: HepG2 cells,

Background Information

CNPY2, also named as MSAP, TMEM4 and ZSIG9, is a positive regulator of neurite outgrowth. It interacts with the ezrin-radixin-moesin (ERM)-like myosin regulatory light chain-interacting protein (MIR), and the two proteins are colocalized in cell lines and in primary neurons. CNPY2 can stabilize MRLC and thus bring about an increase in neurite outgrowth. (PMID: 12826659) This antibody is specific to CNPY2.

Notable Publications

| Author | Pubmed ID | Journal | Application |
|---------------|-----------|---------------------|-------------|
| Feng Hong | 28869608 | Nat Struct Mol Biol | WB |
| Lina Zhao | 31625681 | J Cell Mol Med | WB |
| Chun-Ling Guo | 36422742 | Apoptosis | WB,IF |

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3 $\,$

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




mouse pancreas tissue were subjected to SDS PAGE followed by western blot with 14635-1-AP (CNPY2,MSAP antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 14635-1-AP (CNPY2, MSAP antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




IP result of anti-CNPY2, MSAP (IP:14635-1-AP, 3ug; Detection:14635-1-AP 1:500) with HepG2 cells lysate 600ug.




Immunohistochemical analysis of paraffinembedded human liver tissue slide using 14635-1-AP (CNPY2, MSAP antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Sodium Citrate buffer (pH 6.0).




Immunofluorescent analysis of HepG2 cells using 14635-1-AP (CNPY2,MSAP antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Anti-Human CNPY2, MSAP (14635-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



WB result of CNPY2, MSAP antibody (14635-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CNPY2, MSAP transfected HepG2 cells.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 14635-1-AP (CNPY2, MSAP antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Sodium Citrate buffer (pH 6.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CNPY2, MSAP antibody (14635-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).