For Research Use Only

## NKRF Polyclonal antibody

Catalog Number:14693-1-AP

Featured Product

4 Publications





Basic Information	Catalog Number: GenBank Accession Number: 14693-1-AP BC068514		lumber:	Purification Method: Antigen affinity purification	
	Size: GeneID (NCBI):   450 μg/ml 55922			Recommended Dilutions:	
				WB 1:5000-1:50000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate	
	Source: Rabbit	UNIPROT ID: 015226			
	Isotype: Full Name: IgG NFKB repressir		Dr		
	Immunogen Catalog Number: AG6252	Calculated MW: 78 kDa			
		Observed MW: 100 kDa			
Applications	Tested Applications:	Positive Controls:			
	IP, WB, ELISA		WB: HeLa cells, HEK-293 cells, HepG2 cells, Jurkat		
	Cited Applications: WB, IF, CoIP, ChIP		cells		
	IP : HeLa cells, Species Specificity: human, mouse, rat				
	Cited Species: human, mouse				
Background Information	NKRF, also named as Protein ITBA4 or Transcription factor NRF, is a 690 amino acid protein, which contains one G- patch domain and one R3H domain. NKRF is Widely and constitutively expressed, with lower level in colon, peripheral blood lymphocytes, lung and kidney. NKRF interacts with a specific negative regulatory element (NRE) 5'- AATTCCTCTGA-3' to mediate transcriptional repression of certain NK-kappa-B responsive genes. NKRF is involved in the constitutive silencing of the IFN- $\beta$ promoter, independently of the virus-induced signals, and in the inhibition of the basal and cytokine-induced iNOS promoter activity.				
Notable Publications	Author	Pubmed ID Journ	al	Application	
	Xi Jin	32393270 Mol C	ancer	WB	
	Huang Lisong L	23808420 Immu	inology	WB	
	Yamin Liu	38612387 Int J M	1ol Sci	WB	
Storage	Storage: Store at -20°C. Stable for one yea Storage Buffer:	ar after shipment.			


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data







Various lysates were subjected to SDS PAGE followed by western blot with 14693-1-AP (NKRF antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours.

WB result of NKRF antibody (14693-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-NKRF transfected HEK-293 cells. IP result of anti-NKRF (IP:14693-1-AP, 4ug; Detection:14693-1-AP 1:10000) with HeLa cells lysate 840 ug.