For Research Use Only

PPP2R3A Polyclonal antibody

Catalog Number: 14720-1-AP

2 Publications



Basic Information

 Catalog Number:
 GenBank Accession

 14720-1-AP
 BC065531

 Size:
 GeneID (NCBI):

 153 μg/ml
 5523

 Source:
 UNIPROT ID:

 Rabbit
 Q06190

Full Name: protein phosphatase 2 (formerly 2A), regulatory subunit B", alpha

Immunogen Catalog Number: regulatory subul AG6450 Calculated MW:

130 kDa

Observed MW:

130 kDa

Applications

Tested Applications: IHC, WB,ELISA
Cited Applications:

WB, IF

Isotype:

Species Specificity: human Cited Species:

human

Note-IHC: suggested antigen retrieval with

retrieval may be performed with citrate buffer pH 6.0

GenBank Accession Number: Purification Method:
BC065531 Antigen affinity purification
GenelD (NCBI): Recommended Dilutions:
5523 WB 1:500-1:2000
IHC 1:20-1:200

Positive Controls:

WB: PC-3 cells, MCF-7 cells

IHC: human skin tissue, human brain tissue, human kidney tissue, human lung tissue, human ovary tissue, human placenta tissue, human spleen tissue, human

testis tissue

Background Information

PPP2R3A(Serine/threonine-protein phosphatase 2A regulatory subunit B" subunit alpha), also named as PPP2R3, has 2 isoforms produced by alternative splicing named as PR130 and PR72. The protein contains stretches of basic residues that may be putative nuclear localization signals, and a cluster of proline residues in an N-terminal region common to both isoforms that the isoform PR130 but not PR72 is variably expressed in a number of tumor cell lines. The PR72 and PR130 cDNAs encode 72 and 130 kDa proteins when expressed in rabbit reticulocyte lysates. (PMID:8392071).

Notable Publications

Author	Pubmed ID	Journal	Application
Xiao-Nian Zhu	24412756	Cell Signal	WB
Xiaoni Shi	35933397	Cell Biosci	WB,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


TE buffer pH 9.0; (*) Alternatively, antigen

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



PC-3 cells were subjected to SDS PAGE followed by western blot with 14720-1-AP (PPP2R3A antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human skin using 14720-1-AP (PPP2R3A antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human skin using 14720-1-AP (PPP2R3A antibody) at dilution of 1:100 (under 40x lens).