

For Research Use Only

Noggin Polyclonal antibody

Catalog Number: 14772-1-AP **2 Publications**



Basic Information

Catalog Number: 14772-1-AP	GenBank Accession Number: BC034027	Purification Method: Antigen affinity purification
Size: 500 µg/ml	GeneID (NCBI): 9241	Recommended Dilutions: WB 1:500-1:3000 IHC 1:20-1:200
Source: Rabbit	UNIPROT ID: Q13253	
Isotype: IgG	Full Name: noggin	
Immunogen Catalog Number: AG6586	Calculated MW: 26 kDa	
	Observed MW: 26 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls: WB : HeLa cells, IHC : human gliomas tissue,
Cited Applications: WB, IHC	
Species Specificity: human, mouse, rat	
Cited Species: mouse, hamster	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

Noggin is an extracellular polypeptide acting as an antagonist of bone morphogenetic proteins (BMPs) regulating embryonal development. Noggin inhibits activity of BMP-2, -4, -7, -13, and -14. Noggin is present extracellularly in the matrix or retained at the cell surface via interaction with heparin sulfate proteoglycans. In early development stages, Noggin is produced by the Spemann organizer, allowing dorsal-ventral patterning of BMPs (PMID: 8752214). Subsequently, Noggin is expressed by the notochord regulating BMP-4 signaling in neurogenesis. Additionally, Noggin is present during development in the dermal papilla, connective tissue of the hair follicle, lens, retina, and pericardial mesenchyme, as well as in the mesoderm lineage regulating development of the bone, cartilage, and muscles.

Notable Publications

Author	Pubmed ID	Journal	Application
Chuanhui Yin	32949110	Birth Defects Res	IHC
Shu-Fen Li	29415997	Cell Death Dis	WB

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



HeLa cells were subjected to SDS PAGE followed by western blot with 14772-1-AP (Noggin antibody at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human gliomas using 14772-1-AP (Noggin antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human gliomas using 14772-1-AP (Noggin antibody) at dilution of 1:50 (under 40x lens).