### For Research Use Only

# PEX16 Polyclonal antibody

Catalog Number: 14816-1-AP

Featured Product

14 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 14816-1-AP BC000467

Size: GeneID (NCBI): 9409

Source: UNIPROT ID:

Source: UNIPROT II
Rabbit Q9Y5Y5
Isotype: Full Name:

IgG peroxisomal biogenesis factor 16

Immunogen Catalog Number: Calculated MW:

AG6566 39 kDa

Observed MW: 38 kDa Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:2000

IF/ICC 1:20-1:200

**Applications** 

Tested Applications: WB, IF/ICC, ELISA Cited Applications: WB, IHC, IF Species Specificity: human, mouse, rat

**Cited Species:** 

human, mouse, arabidopsis

Positive Controls:

WB: human liver tissue, HepG2 cells

IF/ICC: HepG2 cells,

## **Background Information**

Peroxins (PEXs) are proteins that are essential for the assembly of functional peroxisomes. Peroxin 16, also known as PEX16 or Peroxisomal biogenesis factor 16, is a 336 amino acid integral membrane protein that has a critical role in the biogenesis of peroxisomes. PEX16 together with PEX3 and PEX19 are specifically involved in peroxisomal membrane protein (PMP) import. Defects in the gene encoding Peroxin 16 are the cause of multiple peroxisome-related disorders, including Zellweger syndrome (ZWS), neonatal adrenoleukodystrophy (NALD), infantile Refsum disease (IRD), classical rhizomelic chondrodysplasia punctata (RCDP) and peroxisome biogenesis disorder complementation group 9 (PBD-CG9).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Lingjuan Piao	31530170	Antioxid Redox Signal	WB
Ming-Ying Ling	36326546	Arch Biochem Biophys	IHC
Katharina M Walter	25440060	Cell Metab	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.


Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**







human liver tissue were subjected to SDS PAGE followed by western blot with 14816-1-AP (PEX16 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immun of luorescent analysis of HepG2 cells using 14816-1-AP (PEX16 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Affini Pure Goat Anti-Rabbit IgG(H+L).