

CXCR7-Specific Polyclonal antibody

Catalog Number: 14840-1-AP

Basic Information

Catalog Number:

14840-1-AP

Size:

170 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_020311

GeneID (NCBI):

57007

UNIPROT ID:

P25106

Full Name:

chemokine (C-X-C motif) receptor 7

Calculated MW:

41 kDa

Observed MW:

50 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:200-1:1000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

Applications

Tested Applications:

IHC, IP, WB, ELISA

Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with **TE buffer pH 9.0; (*)** Alternatively, antigen retrieval may be performed with **citrate buffer pH 6.0**

Positive Controls:

WB : K-562 cells, C6 cells, mouse brain tissue

IP : K-562 cells,

IHC : mouse brain tissue,

Background Information

CXCR7 (C-X-C chemokine receptor type 7), also known as RDC1, is a member of the G-protein coupled receptor family. CXCR7 can bind the chemokines CXCL11 and CXCL12 with high affinity, and it also acts as coreceptor with CXCR4 for a restricted number of HIV isolates. Expression of CXCR7 has been associated with cardiac development as well as with tumor progression. This antibody is specific to CXCR7.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




K-562 cells were subjected to SDS PAGE followed by western blot with 14840-1-AP (CXCR7-Specific antibody) at dilution of 1:100 incubated at room temperature for 1.5 hours.



IP result of anti-CXCR7-Specific (IP:14840-1-AP, 4ug; Detection:14840-1-AP 1:500) with K-562 cells lysate 3200ug.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 14840-1-AP (CXCR7-Specific Antibody) at dilution of 1:100 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 14840-1-AP (CXCR7-Specific Antibody) at dilution of 1:100 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).