

For Research Use Only

SYK Polyclonal antibody

Catalog Number: 14858-1-AP

Featured Product

4 Publications



Basic Information

Catalog Number:

14858-1-AP

Size:

800 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG6657

GenBank Accession Number:

BC001645

GeneID (NCBI):

6850

ENSEMBL Gene ID:

ENSG00000165025

UNIPROT ID:

P43405

Full Name:

spleen tyrosine kinase

Calculated MW:

72 kDa

Observed MW:

66-72 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IHC 1:400-1:1600

Applications

Tested Applications:

IHC, WB, ELISA

Cited Applications:

WB, IHC

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Positive Controls:

WB: Daudi cells, Raji cells, Ramos cells

IHC: human liver cancer tissue, human appendicitis tissue, human spleen tissue, human kidney tissue, human ovary tissue

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

SYK belongs to the protein kinase superfamily, Tyr protein kinase family and SYK/ZAP-70 subfamily. It is positive effector of BCR-stimulated responses. SYK couples the B-cell antigen receptor (BCR) to the mobilization of calcium ion either through a phosphoinositide 3-kinase-dependent pathway, when not phosphorylated on tyrosines of the linker region, or through a phospholipase C-gamma-dependent pathway. It phosphorylates USP25 and regulates its intracellular levels. SYK also promotes liver fibrosis by activation of hepatic stellate cells and acts as a biomarker for human hepatocellular carcinoma. (PMID: 29537660) SYK can be detected 66-72 kDa in human and mouse (PMID: 37661351).

Notable Publications

Author	Pubmed ID	Journal	Application
Le Pei	36359082	Animals (Basel)	WB
Quanwei Zhou	35910221	Front Genet	IHC
Jing Zhang	38320352	Int Immunopharmacol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 14858-1-AP (SYK antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



WB result of SYK antibody (14858-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-SYK transfected Raji cells.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 14858-1-AP (SYK antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).