For Research Use Only

QDPR Polyclonal antibody

Catalog Number: 14908-1-AP

3 Publications



Purification Method:

WB 1:500-1:2000

protein lysate

IHC 1:20-1:200

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 14908-1-AP BC000576
Size: GeneID (NCBI): 5860
Source: UNIPROT ID: Rabbit P09417

Isotype: Full Name:
IgG quinoid dihydropteridine reductase

9- 4------

28-30 kDa

Immunogen Catalog Number:Calculated MW:AG670526 kDa

Observed MW:

Applications

Tested Applications: IHC, IP, WB, ELISA Cited Applications:

WB

Species Specificity: human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: mouse liver tissue, human liver tissue

IP: mouse liver tissue,

IHC: human ovary tumor tissue,

Background Information

Dihydropteridine reductase (QDPR), also named as DHPR and HDHPR, is an essential enzyme in the hydroxylating system of the aromatic amino acids, since it catalyses the regeneration of tetrahydrobiopterin (BH4), the natural cofactor of phenylalanine, tyrosine, and tryptophan hydroxylases, from the quininoid-dihydrobiopterin produced in these coupled reactions(PMID:8326489). The QDPR protein is active as a dimer, with a subunit Mr of 26 kDa(PMID:7627180). This protein belongs to the short-chain dehydrogenases/reductases (SDR) family. Defects in QDPR are the cause of BH4-deficient hyperphenylalaninemia type C (HPABH4C)(PMID:11153907).

Notable Publications

Author	Pubmed ID	Journal	Application
Natasha L Pacheco	29090078	Mol Autism	WB
Zhiguo Li	34643893	Genes Genomics	WB
Kalina Wiatr	34220448	Front Mol Neurosci	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




mouse liver tissue were subjected to SDS PAGE followed by western blot with 14908-1-AP (QDPR antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human ovary tumor using 14908-1-AP (QDPR antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human ovary tumor using 14908-1-AP (QDPR antibody) at dilution of 1:100 (under 40x lens).



IP result of anti-QDPR (IP:14908-1-AP, 4ug; Detection:14908-1-AP 1:1000) with mouse liver tissue lysate 4000ug.