For Research Use Only

ZAK Polyclonal antibody

Catalog Number: 14945-1-AP

Featured Product

10 Publications



Basic Information

Catalog Number: GenBank Accession Number: 14945-1-AP BC001401 GeneID (NCBI): Size: 400 μg/ml 51776 **UNIPROT ID:** Source: Rabbit Q9NYL2 Full Name:

WB 1:500-1:1500 IHC 1:50-1:500 IF/ICC 1:200-1:800

Purification Method:

Antigen affinity purification

Recommended Dilutions:

Isotype:

sterile alpha motif and leucine zipper containing kinase AZK

Immunogen Catalog Number:

AG6747 Calculated MW:

> 91 kDa Observed MW: 52 kDa

Applications

Tested Applications: WB, IF, IHC, ELISA Cited Applications:

Species Specificity: human, mouse, rat **Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse skeletal muscle tissue, human heart tissue, HepG2 cells, human skeletal muscle tissue

IHC: human liver cancer tissue, human placenta tissue

IF/ICC: HepG2 cells,

Background Information

ZAK(sterile-alpha motif and leucine zipper containing kinase AZK) is also named as MLTK, MAPKKK, mlklak, MLK7, AZK, MLT, MRK, HCCS-4 and belongs to the MAPKKK family. It is a mitogen-activated protein kinase kinase kinase (MAP3K) that activates the stress-activated protein kinase/c-jun N-terminal kinase pathway and activates NFkappaB. ZAK contributes to regulation of DNA damage checkpoints through a p38gamma-independent pathway. This protein has 3 isoforms produced by alternative splicing with the MW of 91 kDa, 51 kDa and 35 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Goda Snieckute	36384144	Cell Metab	WB
Yan Wang	33667382	Immunity	WB
Joana Silva	35918345	Nat Commun	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




mouse skeletal muscle lysates were subjected to SDS PAGE followed by western blot with 14945-1-AP (ZAK antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 14945-1-AP (ZAK antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 14945-1-AP (ZAK antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using ZAK antibody (14945-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L).