For Research Use Only

UQCRQ Polyclonal antibody

Catalog Number: 14975-1-AP

Featured Product

10 Publications



Basic Information

Catalog Number: GenBank Accession Number: 14975-1-AP BC001390 GeneID (NCBI): Size: 650 µg/ml 27089 **UNIPROT ID:** Source: Rabbit 014949 Full Name: Isotype:

IgG ubiquinol-cytochrome c reductase, complex III subunit VII, 9.5kDa Immunogen Catalog Number:

AG6886 Calculated MW:

10 kDa Observed MW: 10 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:3000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF/ICC 1:50-1:500

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

Species Specificity: human, mouse, rat **Cited Species:** human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HepG2 cells, human liver tissue

IP: HepG2 cells,

IHC: human liver cancer tissue,

IF/ICC: HepG2 cells,

Background Information

UQCRQ(Cytochrome b-c1 complex subunit 8) belongs to the UQCRQ/QCR8 family. UQCRQ is a ubiquinone-binding $protein\ localized\ to\ the\ cytochrome\ bc1\ region\ of\ the\ mitochondrial\ respiratory\ chain.\ The\ deduced\ 110-amino\ acid$ protein has a relatively hydrophobic and basic N-terminal region, an aspartic acid-rich middle region, and a glutamic acid- and lysine-rich C-terminal region.

Notable Publications

| Author | Pubmed ID | Journal | Application |
|-------------------|-----------|----------------|-------------|
| Flora Cozzolino | 32886703 | PLoS One | WB |
| Isabel Carrascoso | 34884582 | Int J Mol Sci | WB |
| Yuehong Wang | 28518143 | Cell Death Dis | WB |

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




HepG2 cells were subjected to SDS PAGE followed by western blot with 14975-1-AP (UQCRQ antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




IP result of anti-UQCRQ (IP:14975-1-AP, 4ug; Detection:14975-1-AP 1:1000) with HepG2 cells lysate 3360 ug.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 14975-1-AP (UQCRQ antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 14975-1-AP (UQCRQ antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using UQCRQ antibody (14975-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).