For Research Use Only

HLA-E Polyclonal antibody

Catalog Number: 15008-1-AP Feat

Featured Product



Purification Method:

WB 1:2000-1:10000

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: 15008-1-AP

Concentration: 500 ug/ml

Source: Rabbit Isotype:

Immunogen Catalog Number:

mmunogen Catalog Numbe

AG6703

....

Tested Applications: WB, ELISA

Species Specificity:

human

GenBank Accession Number:

BC002578 GeneID (NCBI):

UNIPROT ID: P13747 Full Name:

major histocompatibility complex,

class I, E

Calculated MW: 40 kDa

Observed MW: 40-50 kDa

Positive Controls:

WB: HUVEC cells, THP-1 cells, Raji cells

Background Information

Applications

Human major histocompatibility complex (MHC) antigens, also referred to as human leukocyte antigens (HLA), are encoded by genes located on the short arm of chromosome 6 (6p21.3). There are two classes of HLA antigens: class I and class II. This class I molecules are membrane glycoproteins composed of a heavy (alpha) chain which is encoded by a HLA class I gene, and β 2-microglobulin light (beta) chain. The most extensively characterized members of the HLA class I gene family are the genes encoding the major transplantation antigenes, HLA-A, B and C. HLA-E is a non-classical MHC class I molecule. HLA-E is frequently overexpressed in tumor diseases, transplants and virus-infected cells and represents an immunomodulatory molecule by binding to the receptors CD94/NKG2A, -B and -C on NK and T cells. Due to its immune suppressive features HLA-E expression might represent an important mechanism of tumors to escape immune surveillance.(PMID: 667938; 3375250; 2249951; 27589686)


Storage

Storage


Store at -20°C. Stable for one year after shipment. Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol Aliquoting is unnecessary for -20°C storage

Selected Validation Data



WB result of HLA-E antibody (15008-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-HLA-E transfected THP-1 cells.



Various lysates were subjected to SDS PAGE followed by western blot with 15008-1-AP (HLA-E antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.