

For Research Use Only

CITED2 Polyclonal antibody

Catalog Number: 15021-1-AP

1 Publications



Basic Information

Catalog Number:

15021-1-AP

Size:

173 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG7078

GenBank Accession Number:

BC004377

GeneID (NCBI):

10370

UNIPROT ID:

Q99967

Full Name:

Cbp/p300-interacting transactivator, with Glu/Asp-rich carboxy-terminal domain, 2

Calculated MW:

28 kDa

Observed MW:

36 kDa, 27 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IHC 1:50-1:500

IF/ICC 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse, rat

Cited Species:

mouse

Positive Controls:

WB : SH-SY5Y cells, mouse brain tissue

IHC : mouse brain tissue, mouse ovary tissue

IF/ICC : SH-SY5Y cells,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

CITED2, also named as P35srj, is a cAMP-responsive element-binding protein (CBP)/p300-transactivator, with an ED(Glu/Asp)-rich C-terminal domain that acts as an important modulator in the development of the heart [PMID:10552932]. There are two isoforms of CITED2 protein and the calculated molecular weight of them are 24kDa and 28 kDa. The 28 kDa isoform having modification site may migrate to 35 kDa and 24 kDa isoform having modification site may migrate to 27 kDa (PMID: 23082118). It is proposed to be a negative regulator of HIF-1 α through its competitive binding to the CH1 domain of CBP/p300, which mediated signaling and to a decrease in HIF-1-responsive genes such as VEGF [PMID:9887100]. In addition, it is also a co-activator of the transcriptional activity of the TFAP2 isoform, which is an extensively expressed nuclear protein with functions in embryogenesis, inflammation, and stress responses[PMID:22735262].

Notable Publications

Author	Pubmed ID	Journal	Application
Yan Jiang	28013219	Toxicol Sci	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




SH-SY5Y cells were subjected to SDS PAGE followed by western blot with 15021-1-AP (CITED2 antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (10% Formaldehyde) fixed SH-SY5Y cells using 15021-1-AP (CITED2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 15021-1-AP (CITED2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 15021-1-AP (CITED2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse ovary tissue slide using 15021-1-AP (CITED2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).