For Research Use Only

SMOX Polyclonal antibody

Catalog Number:15052-1-AP

Featured Product






Basic Information	Catalog Number: 15052-1-AP	GenBank Accessio BC000669	on Number:	Purification Method: Antigen affinity purification	
	Concentration: 300 μg/ml	Genel D (NCBI): 54498		Recommended Dilutions: WB 1:2000-1:10000	
	Source: Rabbit	UNIPROT ID: Q9NWM0		IHC 1:50-1:500	
	Isotype: IgG	Full Name: spermine oxidase			
	Immunogen Catalog Number: AG7035	Calculated MW: 62 kDa			
		Observed MW: 56-65 kDa			
Applications	Tested Applications:		Positive Con	trols:	
	WB, IHC, ELISA Cited Applications:		WB : A549 cel PC-3 cells	WB : A549 cells, PC-13 cells, MCF-7 cells, HeLa cells PC-3 cells	
				C : human prostate cancer tissue, human stomach ncer tissue, human gliomas tissue	
	Cited Species: human, mouse, rat				
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	SMOX(Spermine oxidase) is also named as SMO, PAO-1 and PAOh1. It belongs to the class of polyamine oxidases (PAOs).The flavoprotein SMO, which specifically oxidizes Spm, is one of the key regulatory enzymes involved in polyamine homeostasis in animal cells (PMID:14764092). It has 6 isoforms produced by alternative splicing with the MW of 61 kDa, 56 kDa, 21 kDa, 60 kDa, 43 kDa and 65 kDa, respectively.				
Notable Publications	Author	Pubmed ID Jo	ournal	Application	
	Takeshi Uemura	32995657 He	eliyon	WB	
	Anna Bianchi-Smiraglia	30198908 J (Clin Invest	WB	
	Bin Xiang	36353478 Fr	ont Pharmacol	IHC	


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




PC-13 cells were subjected to SDS PAGE followed by western blot with 15052-1-AP (SMOX antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 15052-1-AP (SMOX antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 15052-1-AP (SMOX antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 15052-1-AP (SMOX antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 15052-1-AP (SMOX antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 15052-1-AP (SMOX antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).