# For Research Use Only

# TPPP3 Polyclonal antibody

Catalog Number: 15057-1-AP

**Featured Product** 

**3 Publications** 



**Purification Method:** 

WB 1:200-1:1000

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

**Applications** 

Catalog Number: 15057-1-AP Size:

800  $\mu$  g/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG7040

**Tested Applications:** 

WB, ELISA Cited Applications:

Species Specificity: human, mouse, rat **Cited Species:** 

human, mouse

GenBank Accession Number:

BC000691 GeneID (NCBI): 51673 **UNIPROT ID:** Q9BW30

tubulin polymerization-promoting

protein family member 3

Calculated MW: 19 kDa

Observed MW: 19-20 kDa

Full Name:

Positive Controls:

WB: PC-3 cells, HEK-293 cells

# **Background Information**

TPPP3 (tubulin polymerization promoting protein 3), also named as TPPP/p20, is a new member of the TPPP family which binds tubulin and induces tubulin polymerization and microtubule bundling. TPPP3 has been shown to play an analysis of the property ofimportant role in cell proliferation and may be implicated in tumorigenesis and metastasis (19633818). TPPP3 was also identified as specific marker for the connective tissues that envelope tendons due to its distinct expression in the developing tendon sheath (19235716). This antibody detects the 19-20 kDa TPPP3 in HeLa and PC-3 cells. The non-specific bands between 40 and 60 kDa were also occasionally observed.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Zhen Xiong	35320705	Immunity	
Mengqi Chu	33509274	Stem Cell Res Ther	WB
Zhen Huang	37483625	Front Immunol	IF


### Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

# Selected Validation Data





WB result of anti-TPPP3 (15057-1-AP, 1:500) with HEK-293 cells (siRNA transfected with) by Dr. Yupeng Chen. (Negative control or TPPP3 specific siRNAs were transfected into HEK-293 cells with the final concentration of 20 nM. Forty eight hours after transfection, cells were lysed with RIPA buffer and subjected to SDS PAGE).

PC-3 cells were subjected to SDS PAGE followed by western blot with 15057-1-AP (TPPP3 antibody) at dilution of 1:100 incubated at room temperature for 1.5 hours.