## For Research Use Only

# GSTT2 Polyclonal antibody

Catalog Number: 15063-1-AP

1 Publications



**Basic Information** 

**Applications** 

Catalog Number: 15063-1-AP Size: 450 µg/ml

Source: Rabbit Isotype:

Immunogen Catalog Number:

AG7066

27 kDa Observed MW: 28 kDa

**Tested Applications:** 

WB, IHC, IF/ICC, ELISA Cited Applications:

WB

Species Specificity: human, mouse, rat **Cited Species:** 

mouse Note-IHC: suggested antigen retrieval with

TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

**Purification Method:** 

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000 IHC 1:100-1:400 IF/ICC 1:20-1:200

Positive Controls:

GenBank Accession Number:

glutathione S-transferase theta 2

BC002415

2953

POCG29 Full Name:

GeneID (NCBI):

**UNIPROT ID:** 

Calculated MW:

WB: rat liver tissue, LO2 cells, mouse liver tissue,

mouse lung tissue

IHC: human prostate cancer tissue,

IF/ICC: LO2 cells,

# **Background Information**

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Cheng-Niu Wang	32987309	Bioorg Chem	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


### **Selected Validation Data**




rat liver tissue were subjected to SDS PAGE followed by western blot with 15063-1-AP (GSTT2 Antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 15063-1-AP (GSTT2 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 15063-1-AP (GSTT2 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of LO2 cells using 15063-1-AP (GSTT2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).