For Research Use Only

POLDIP2 Polyclonal antibody

Catalog Number: 15080-1-AP

3 Publications



Basic Information

Catalog Number: 15080-1-AP Size: 273 µg/ml

Rabbit Isotype:

Immunogen Catalog Number:

Immunogen Catalog Number:

AG7154

Source:

GenBank Accession Number:

BC000655

GeneID (NCBI):
26073

UNIPROT ID:
Q9Y257

Full Name:

polymerase (DNA-directed), delta

interacting protein 2
Calculated MW:
42 kDa

Observed MW: 38 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:500-1:3000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IF 1:10-1:100

Applications

Tested Applications: IF/ICC, IP, WB, ELISA Cited Applications:

WB, IF

Species Specificity: human, mouse, rat Cited Species: human, mouse Positive Controls:

WB: HeLa cells, HepG2 cells, Jurkat cells, K-562 cells

IP : HeLa cells,
IF : Hela cells,

Background Information

POLDIP2(Polymerase delta-interacting protein 2) is also named as PDIP38, POLD4. It is as a unique positive regulator of Nox4 via its association with p22phox and together with Nox4, has profound effects on Rhodependent cytoskeletal remodeling. It is also as a regulator of cell division. It consists of 368 amino acids and has a predicted molecular mass of 42 kDa, with a potential signal peptide cleavage site after the first N-terminal 48 residues, which would result in a protein of 37 kDa(PMID:19574552). Endogenous human and rat POLDIP2 localized to the nucleus, plasma membrane, and cytoplasm(PMID:17623671).

Notable Publications

Author	Pubmed ID	Journal	Application
Agnes Wong	23989611	Cell Cycle	IF
Jana Key	34943861	Cells	WB
Maga Giovanni G	24191025	Proc Natl Acad Sci U S A	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



HeLa cells were subjected to SDS PAGE followed by western blot with 15080-1-AP (POLDIP2 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of Hela cells, using POLDIP2 antibody 15080-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



IP result of anti-POLDIP2 (IP:15080-1-AP, 4ug; Detection:15080-1-AP 1:1000) with HeLa cells lysate 1600ug.