For Research Use Only

IL-10RB Polyclonal antibody Catalog Number:15102-1-AP 4 Publications




Basic Information	Catalog Number: 15102-1-AP	GenBank Accession Num BC001903	ber: Purification Meth Antigen affinity	
	Size:	GenelD (NCBI):	Recommended D	vilutions:
	450 µg/ml	3588	WB 1:300-1:1000	
	Source:	UNIPROT ID:		
	Rabbit	Q08334		
	Isotype: Full Name: IgG interleukin 10 receptor, beta		oeta	
	Immunogen Catalog Number:	Calculated MW:		
	AG7445	37 kDa		
		Observed MW:		
		42-45 kDa		
Applications	Tested Applications:	Р	ositive Controls:	
	WB, ELISA		WB : rat skeletal muscle tissue, Jurkat cells, mouse	
	Cited Applications: WB	Cited Applications: skeletal muscle tissue WB		
	Species Specificity:			
	human, mouse, rat			
	Cited Species:			
	human, mouse			
Background Information	Interleukin-10 (IL-10) is an anti-inflammatory cytokine that plays a critical role in maintaining immune system balance. IL-10 signals through a receptor complex consisting of two molecules of the IL-10R alpha-chain (IL-10RA) and two molecules of the accessory IL-10R beta-chain (IL-10RB) (PMID: 22236434). IL-10RA and IL-10RB are member of the type II cytokine receptor family. IL-10RB, also known as IL-10R2 or CDw210b, is the signaling subunit of the IL-10R complex and is constitutively expressed in a variety of cell types (PMID: 24507158). Binding of IL-10 to the IL-10R complex leads to JAK1- and TYK2-mediated phosphorylation of STAT3 (PMID: 11244051; 10433356). IL-10RB is also shared by several other cytokines, including IL22, IL-26, IFN- γ, IL-28A/B and IL29 (PMID: 22236434).			
Notable Publications	Author	Pubmed ID Journal		Application
	Yang Che	34015450 Biochim	Biophys Acta Mol Basis Dis	WB
	Shuyue Luo	35811529 J Alzhein	ners Dis	WB
	Yao Jiang	38647150 Adv Heal	thc Mater	WB

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



rat skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 15102-1-AP (IL10RB antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.