

For Research Use Only

# WDR46 Polyclonal antibody

Catalog Number: 15110-1-AP

4 Publications



## Basic Information

### Catalog Number:

15110-1-AP

### Size:

1000 ug/ml

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG7095

### GenBank Accession Number:

BC000388

### GeneID (NCBI):

9277

### UNIPROT ID:

O15213

### Full Name:

WD repeat domain 46

### Calculated MW:

68 kDa

### Observed MW:

70-75 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB 1:2000-1:10000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

## Applications

### Tested Applications:

WB, FC (Intra), IP, ELISA

### Cited Applications:

WB, IF

### Species Specificity:

human, mouse, rat

### Cited Species:

human, mouse

### Positive Controls:

WB : HeLa cells, Jurkat cells, mouse spleen tissue, rat spleen tissue

IP : Jurkat cells,

## Background Information

## Notable Publications

Author	Pubmed ID	Journal	Application
Kwon Tae You	26443847	Genes Dev	WB
Kouko Wada	24754225	Biochem Cell Biol	IF
Hirai Yuya Y	23848194	Genes Cells	WB,IF

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Jurkat cells were subjected to SDS PAGE followed by western blot with 15110-1-AP (WDR46 antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 15110-1-AP (WDR46 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



1x10<sup>6</sup> Jurkat cells were intracellularly stained with 0.25 ug WDR46 Polyclonal antibody (15110-1-AP) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



IP result of anti-WDR46 (IP:15110-1-AP, 4ug; Detection:15110-1-AP 1:8000) with Jurkat cells lysate 2280 ug.



IP result of anti-WDR46 (IP:15110-1-AP, 4ug; Detection:15110-1-AP 1:4000) with Jurkat cells lysate 1400 ug.