## For Research Use Only

# SIRT5 Polyclonal antibody

Catalog Number: 15122-1-AP

Featured Product

51 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 15122-1-AP BC000126

Concentration: GeneID (NCBI): 23408

Source: UNIPROT ID: Rabbit Q9NXA8

Isotype:Full Name:IgGsirtuin (silent

Immunogen Catalog Number: informatio

AG7195

sirtuin (silent mating type information regulation 2 homolog) 5 (S. cerevisiae)

Calculated MW: 34 kDa Observed MW: 33 kDa Purification Method:
Antigen affinity purification
Recommended Dilutions:

WB 1:2000-1:16000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IF/ICC 1:50-1:500

**Applications** 

Tested Applications:
WB, IF/ICC, IP, ELISA
Cited Applications:
WB, IHC, IF, IP, CoIP
Species Specificity:
human, mouse, rat
Cited Species:
human, mouse, rat, bovine

#### **Positive Controls:**

WB: HCT 116 cells, RAW 264.7 cells, mouse heart tissue, mouse liver tissue, rat heart tissue, rat liver

tissue

IP: mouse liver tissue,
IF/ICC: HeLa cells,

# **Background Information**

SIRT5, a member fo sirtuin family that is a mitochondrial matrix NAD(+)-dependent deacetylase and mono-ADP-ribosyltransferases. Sirtuins play important role in various cellular processed, such as aging, gene silencing and metabolism. SIRT5 mainly locates in cytoplasm. It is an efficient protein lysine desuccinylase and demalonylase. During prolonged fasting, it can activate CPS1, the committed and regulated enzyme of the urea cycle, and contribute to the regulation of blood ammonia levels.

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yipeng Du	30279144	EBioMedicine	WB,IF,IHC
Xiaodan Sun	31572453	Front Genet	IHC
Min Liu	28871079	Nat Commun	WB

# Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


# Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 15122-1-AP (SIRT5 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



IP result of anti-SIRT5 (IP:15122-1-AP, 4ug; Detection:15122-1-AP 1:500) with mouse liver tissue lysate 6000ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using 15122-1-AP (SIRT5 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).