For Research Use Only

# NNMT Polyclonal antibody

Catalog Number:15123-1-AP

Featured Product





### **Basic Information**

15123-1-AP Concentration: 500 ug/ml Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG7197

Catalog Number:

GenBank Accession Number: BC000234 GeneID (NCBI): 4837 UNIPROT ID: P40261 Full Name: nicotinamide N-methyltransferase Calculated MW: 30 kDa Observed MW: 30 kDa

#### Purification Method: Antigen affinity purification

#### Recommended Dilutions:

WB 1:1000-1:8000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:400-1:1600 IF/ICC 1:20-1:200

## Applications

Tested Applications: WB, IHC, IF/ICC, FC (Intra), IP, ELISA Cited Applications: WB, IHC, IF, IP Species Specificity:

human, mouse, rat Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### Positive Controls:

WB : LO2 cells, HepG2 cells, mouse liver tissue, rat liver tissue IP : HepG2 cells,

IHC : human stomach cancer tissue,

IF/ICC : HepG2 cells,

### **Background Information**

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Changmei Yang	36312432	ACS Omega	WB
Changqing Wu	36382559	Adv Sci (Weinh)	WB,IHC
Wanfeng Xu	35603729	Bioengineered	WB

#### Storage


#### Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol, pH7.3 Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


### Selected Validation Data




IP result of anti-NNMT (IP:15123-1-AP, 4ug; Detection:15123-1-AP 1:300) with HepG2 cells lysate 2400ug.




Immunofluorescent analysis of (10% Formaldehyde) fixed HepC2 cells using 15123-1-AP (NNMT antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Anti-Human NNMT (15123-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).





Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 15123-1-AP (NNMT antibody) at dilution of 1:800 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

LO2 cells were subjected to SDS PAGE followed by western blot with 15123-1-AP (NNMT antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.