For Research Use Only

HDGF2 Polyclonal antibody

Catalog Number: 15134-1-AP

Featured Product

15 Publications



Purification Method:

IHC: 1:50-1:500 IF/ICC: 1:200-1:800

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number:

15134-1-AP

BC000755

Concentration:

1000 ug/ml

Source:

Rabbit

Q7Z4V5

Isotype:

GenBank Accession Number:

BC000755

GeneID (NCBI):

BC000755

UNIPROT ID:

Q7Z4V5

Full Name:

hepatoma-derived growth factor-

Immunogen Catalog Number: related protein 2
AG7281 Calculated MW:
74 kDa

Applications

Tested Applications: IHC, IF/ICC, ELISA Cited Applications: IF, CoIP, chIP Species Specificity:

Cited Species: human, mouse

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

IHC: human colon tissue,
IF/ICC: HepG2 cells, A431 cells

Background Information

HDGF2, also known as HDGFL2 (Hepatoma-Derived Growth Factor-like 2), is a gene that encodes a member of the hepatoma-derived growth factor (HDGF) family. HDGF2 plays a role in LEDGF/p75-independent HIV-1 replication by determining HIV-1 integration site selection. It is also involved in the binding to methylated histones, which suggests a role in epigenetic regulation.

Notable Publications

Author	Pubmed ID	Journal	Application
Judd F Hultquist	27783955	Cell Rep	WB
Xiao-Feng Tian	27779720	Mol Med Rep	
Lu Liu	29285117	Exp Ther Med	

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3 $\,$

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 15134-1-AP (HRP 2 antibody at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 15134-1-AP (HRP 2 antibody at dilution of 1:200 (under 40x lens).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using HDGF2 antibody (15134-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).