## For Research Use Only

# IFIT3 Polyclonal antibody

Catalog Number: 15201-1-AP

**Featured Product** 

40 Publications

BC004977

GeneID (NCBI):

GenBank Accession Number:



**Basic Information** 

Catalog Number: 15201-1-AP Concentration: 600 µg/ml

Source: UNIPROT ID:
Rabbit 014879
Isotype: Full Name:
IgG IFIT3

Immunogen Catalog Number: Calculated MW:

AG7349 56 kDa

Observed MW: 60 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:2000-1:12000

IHC 1:50-1:500

**Applications** 

Tested Applications: WB, IHC, ELISA

Cited Applications: WB, IHC, IF, ChIP Species Specificity:

human Cited Species:

human, mouse, pig, monkey

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: MKN-45 cells, SGC-7901 cells IHC: human spleen tissue,

# **Background Information**

IFIT3 (IFN-induced protein with tetratricopeptide repeats 3), also known as IFIT-4, ISG-60 or RIG-G, is a 56 kDa IFN induced protein. IFIT3 has been identified to be associated with strong IFN antiviral activity. Upon RNA virus infection, IFIT3 is significantly induced, ectopic expression or knockdown of IFIT3 could, respectively, enhance or impair IFN regulatory factor 3 (IRF3)-mediated gene expression. IFIT3 thus may serve as an important modulator in innate immunity, revealing a new function of the IFIT family proteins.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Zhangtao Jiang	27681138	J Virol	WB
Jiaming Su	36163413	J Med Virol	WB
Liu Xin-Yi XY	21813773	J Immunol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 15201-1-AP (IFIT3 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human spleen tissue slide using 15201-1-AP (IFIT3 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human spleen tissue slide using 15201-1-AP (IFIT3 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).