For Research Use Only

MAEA Polyclonal antibody

Catalog Number: 15238-1-AP

1 Publications



Basic Information

Catalog Number: GenBank Accession Number: 15238-1-AP BC001225
Size: GeneID (NCBI): 350 µg/ml 10296
Source: UNIPROT ID:

Source: UNIPROT II
Rabbit Q7L5Y9
Isotype: Full Name:

IgG macrophage erythroblast attacher

Immunogen Catalog Number: Calculated MW:

AG7365 45 kDa

Observed MW: 42-45 kDa

42-45 kDa

Applications

Tested Applications: IHC, WB, ELISA
Cited Applications:

WB

Species Specificity:

human
Cited Species:
human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HepG2 cells, Jurkat cells, MCF-7 cells, U2OS cells IHC: human lung cancer tissue, human breast cancer

Purification Method:

WB 1:500-1:2000 IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Background Information

MAEA (macrophage erythroblast attacher) is a core component of the CTLH E3 ubiquitin-protein ligase complex. MAEA is required for normal cell proliferation and plays a role in erythroblast enucleation during erythrocyte maturation and in the development of mature macrophages. N-terminus MAEA and full-length MAEA are mostly expressed in nuclear and can be detected at low level in the cytoplasm of some cells, while C-terminus MAEA can be detected in the cytoplasm and nucleoli within the nucleus. MAEA mediates the attachment of erythroid cell to mature macrophages which can inhibit erythroid cell apoptosis. The expression of MAEA may be associated with the type 2 diabetes mellitus. (PubMed: 28955747, 24143168, 29911972, 29911972, 30674470, 9763581).

Notable Publications

Author	Pubmed ID	Journal	Application
Lalit R Patel	37426340	iScience	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 15238-1-AP (MAEA antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 15238-1-AP (MAEA antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).