

## HLA class I ABC Polyclonal antibody

Catalog Number: 15240-1-AP

59 Publications

## Basic Information

## Catalog Number:

15240-1-AP

## Concentration:

900 ug/ml

## Source:

Rabbit

## Isotype:

IgG

## Immunogen Catalog Number:

AG7370

## GenBank Accession Number:

BC003069

## GeneID (NCBI):

3105

## UNIPROT ID:

P04439

## Full Name:

major histocompatibility complex, class I, A

## Calculated MW:

41 kDa

## Observed MW:

44 kDa

## Purification Method:

Antigen affinity purification

## Recommended Dilutions:

WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:5000-1:20000

IF/ICC 1:50-1:500

## Applications

## Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

## Cited Applications:

WB, IHC, IF, IP, CoIP, RIP, Cell treatment

## Species Specificity:

human

## Cited Species:

human

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Positive Controls:

WB : A549 cells, NCI-H1299 cells, Calu-1 cells, HEK-293 cells, HeLa cells, HepG2 cells, L02 cells

IP : HepG2 cells,

IHC : human tonsillitis tissue, human stomach cancer tissue

IF/ICC : HepG2 cells, Raji cells

## Background Information

Human major histocompatibility complex (MHC) antigens, also referred to as human leukocyte antigens (HLA), are encoded by genes located on the short arm of chromosome 6 (6p21.3). There are two classes of HLA antigens: class I (HLA-A, B and C) and class II (HLA-D). This class I molecules are polymorphic membrane glycoproteins composed of a heavy (alpha) chain (44 kDa) which is encoded by a HLA class I gene (HLA-A, B or C), and  $\beta$  2-microglobulin light (beta) chain (12 kDa). They are involved in the presentation of foreign antigens to the immune system. (PMID: 667938; 3375250)

## Notable Publications

Author	Pubmed ID	Journal	Application
Zachary C Dobbin	25209969	Oncotarget	IHC
Tianxiao Xu	34571002	J Invest Dermatol	WB
Rupert L Mayer	36241641	Nat Commun	WB

## Storage

## Storage:

Store at -20°C. Stable for one year after shipment.

## Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 15240-1-AP (HLA class I ABC antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 15240-1-AP (HLA class I ABC antibody) at dilution of 1:10000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 15240-1-AP (HLA class I ABC antibody) at dilution of 1:10000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using HLA class I ABC antibody (15240-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IP result of anti-HLA class I ABC (IP:15240-1-AP, 3ug; Detection:15240-1-AP 1:500) with HepG2 cells lysate 1720ug.



Immunofluorescent analysis of (4% PFA) fixed Raji cells using HLA class I ABC antibody (15240-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).