For Research Use Only

Elastin Polyclonal antibody

Catalog Number: 15257-1-AP 26 Publications



Basic Information

Catalog Number: GenBank Accession Number: 15257-1-AP BC065566

Concentration: GeneID (NCBI): 400 ug/ml 2006

Source: UNIPROT ID:

Rabbit P15502
Isotype: Full Name: lgG elastin

Immunogen Catalog Number: Calculated MW:

AG7179 68 kDa

Observed MW: 60-70 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:1000-1:5000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:500-1:2000

Applications

Tested Applications: WB, IHC, IP, ELISA Cited Applications: WB, IHC, IF Species Specificity:

human Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: human artery tissue,

IP: human placenta tissue,

IHC: human lung tissue, human skin cancer tissue

Background Information

Elastic fibers are an abundant and integral part of many extracellular matrices, in which they provide the elastic properties to tissues such as arterial, lung, and skin. Elastic fibers are consisting of an elastin core surrounded by a mantle of fibrillin-rich microfibrils (PMID: 12082143). Elastin is an extremely durable, insoluble biopolymer formed through the lysine-mediated crosslinking of its soluble precursor tropoelastin, which is an approximately 60-70 kDa protein (PMID: 15837523). Deletions and mutations in the elastin gene (ELN) are associated with supravalvular aortic stenosis (SVAS) and autosomal dominant cutis laxa.

Notable Publications

Author	Pubmed ID	Journal	Application
Yuqi Li	36132982	J Immunol Res	IHC
Avinash Soundararajan	34831087	Cells	WB
Jing Ge	29019707	Am J Respir Cell Mol Biol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Human artery tissue were subjected to SDS PAGE followed by western blot with 15257-1-AP (Elastin antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human lung tissue slide using 15257-1-AP (Elastin antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung tissue slide using 15257-1-AP (Elastin antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-Elastin (IP:15257-1-AP, 4ug; Detection:15257-1-AP 1:300) with human placenta tissue lysate 4000 ug.