

For Research Use Only

# ATF5 Polyclonal antibody

Catalog Number: 15260-1-AP **1 Publications**



## Basic Information

<b>Catalog Number:</b> 15260-1-AP	<b>GenBank Accession Number:</b> BC005174	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 130 µg/ml	<b>GeneID (NCBI):</b> 22809	<b>Recommended Dilutions:</b> WB 1:200-1:1000
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> Q9Y2D1	
<b>Isotype:</b> IgG	<b>Full Name:</b> activating transcription factor 5	
<b>Immunogen Catalog Number:</b> AG7214	<b>Calculated MW:</b> 31 kDa	
	<b>Observed MW:</b> 27-30 kDa	

## Applications

<b>Tested Applications:</b> WB, ELISA	<b>Positive Controls:</b> WB: BxPC-3 cells, A549 cells
<b>Cited Applications:</b> WB	
<b>Species Specificity:</b> human, mouse, rat	
<b>Cited Species:</b> mouse	

## Background Information

ATF5, also named as Activating transcription factor 5, is a 282 amino acid protein, which belongs to the bZIP family. ATF5 is widely expressed with higher expression levels in liver. ATF5 is actively transported to the centrosome and accumulated in the pericentriolar material (PCM) during G1 to M phase via a microtubule-dependent mechanism. During late telophase and cytokinesis, it translocates from the centrosome to the midbody (PMID: 26213385). ATF5 as a transcription factor that either stimulates or represses gene transcription through binding of different DNA regulatory elements such as cAMP response element (CRE) (consensus: 5'-GTGACGT[AC][AG]-3'), ATF5-specific response element (ARE) (consensus: 5'-C[CT]TCT[CT]CCTT[AT]-3') but also the amino acid response element (AARE), present in many viral and cellular promoters. Critically ATF5 is involved, often in a cell type-dependent manner, in cell survival, proliferation, and differentiation (PubMed:10373550, PubMed:15358120, PubMed:21212266, PubMed:20654631).

## Notable Publications

Author	Pubmed ID	Journal	Application
Xu Xu	29698569	Hepatology	WB

## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)

W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



BxPC-3 cells were subjected to SDS PAGE followed by western blot with 15260-1-AP (ATF5 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.