For Research Use Only

## **TNFAIP1** Polyclonal antibody

Catalog Number:15320-1-AP

Featured Product

**3** Publications




Basic Information	Catalog Number: 15320-1-AP	GenBank Accession Number: BC001949		Purification Method: Antigen affinity purification	
	Size: GeneID (NCBI):			Recommended Dilutions:	
	450 µg/ml	7126		WB 1:300-1:1000	
	Source:	Ce. UNIPROTID.		IHC 1:20-1:200 IF/ICC 1:20-1:200	
	Rabbit Q13829				
	Isotype:				
	IgG		tumor necrosis factor, alpha-induced protein 1 (endothelial) Calculated MW:		
	Immunogen Catalog Number: AG0220				
		36 kDa			
		Observed MW:			
		36 kDa			
Applications	Tested Applications:		Positive Controls:		
	WB, IHC, IF/ICC, ELISA		WB : HeLa cells, LO2 cells		
	Cited Applications: WB		IHC : human placenta tissue, human brain tissue		
	Species Specificity:		IF/ICC : HeLa cells,		
	human, mouse, rat				
	Cited Species: human, mouse				
	Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performe buffer pH 6.0	natively, antigen			
Background Information	TNFAIP1 was identified as a human gene activated transcriptionally by tumour necrosis factor alpha in endothel cells. It is encoded by a 3.5-kilobase transcript and its expression is found to be developmentally regulated in a tissue-specific manner. The open reading frame predicts a 316-amino acid sequence rich in charged residues, particularly at the carboxyl terminus.				
Notable Publications	Author	Pubmed ID	Journal	Application	
	Junzhi Yi	32335808	Neurotox Res	WB	
	Wei Wang	38102610	Cell Commun Signa	l WB	
	Zhang Xiaoting X	24179536	Oncol Lett	WB	


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 15320-1-AP (TNFAIP1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human placenta using 15320-1-AP (TNFAIP1 antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human placenta using 15320-1-AP (TNFAIP1 antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of HeLa cells using 15320-1-AP (TNFAIP1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).