For Research Use Only

RPL7A Polyclonal antibody

Catalog Number:15340-1-AP 9 Publications




Basic Information	Catalog Number: 15340-1-AP	GenBank A BC005128	GenBank Accession Number: BC005128		Purification Method: Antigen affinity purification	
	Size:	GenelD (NC	:BI):	Recommended	•	
	450 µg/ml	6130		WB 1:500-1:2000		
	Source:	UNIPROT II	UNIPROT ID: P62424		IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500	
	Rabbit					
	Isotype: IgG	21 C		IF/ICC 1:20-1:200		
	Immunogen Catalog Number: AG7543		Calculated MW: 266 aa, 30 kDa Observed MW: 30-32 kDa			
Applications	Tested Applications:					
	WB, IHC, IF/ICC, IP, ELISA Cited Applications:			MCF7 cells, MCF-7 cells, mouse kidney tissue, se liver tissue, rat kidney tissue		
	WB, IF, IP Species Specificity:		IP : mouse	IP : mouse kidney tissue,		
	human, mouse, rat		IHC : huma	man breast cancer tissue,		
	Cited Species: IF/ICC : HepG			epG2 cells,		
	human, mouse					
	Note-IHC: suggested antigu TE buffer pH 9.0; (*) Altern retrieval may be performe buffer pH 6.0	atively, antig				
Background Information	Ribosomal proteins are a major of component of the 60S large ribos that occur via interaction with hur inhibit the activities of the two n	omal subunit, ha man thyroid hor	s additional functions none receptor (THR) ar	involved in cell grownd retinoic acid recept	wth and differentiation	
Notable Publications	Author	Pubmed ID	Journal		Application	
	Suman Rimal	34663454	Acta Neuropathol Commun		WB	
	Takashi Mino	26000482	Cell		WB, IF	
		31072819	J Exp Med		14/D	
	Hiroki Tanaka	510/2017	5 Expiried		WB	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 15340-1-AP (RPL7A antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



MCF7 cells were subjected to SDS PAGE followed by western blot with 15340-1-AP (RPL7A antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HepG2 cells, using RPL7A antibody 15340-1-AP at 1:50 dilution and Rhodamine-labeled goat anti-rabbit IgG (red). Blue pseudocolor = DAPI (fluorescent DNA dye).



IP result of anti-RPL7A (IP:15340-1-AP, 4ug; Detection:15340-1-AP 1:1000) with mouse kidney tissue lysate 4000ug.