

For Research Use Only

# DHRS3 Polyclonal antibody

Catalog Number: 15393-1-AP

Featured Product

21 Publications



## Basic Information

### Catalog Number:

15393-1-AP

### Size:

600 µg/ml

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG7589

### GenBank Accession Number:

BC002730

### GeneID (NCBI):

9249

### UNIPROT ID:

O75911

### Full Name:

dehydrogenase/reductase (SDR family) member 3

### Calculated MW:

34 kDa

### Observed MW:

30-34 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB 1:500-1:1000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF/ICC 1:200-1:800

## Applications

### Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

### Cited Applications:

WB, IHC, IF

### Species Specificity:

human, mouse, rat

### Cited Species:

human, mouse, insect

### Positive Controls:

WB : A375 cells, mouse liver tissue, rat liver tissue, mouse kidney tissue

IP : A375 cells,

IHC : mouse liver tissue,

IF/ICC : HepG2 cells,

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Background Information

DHRS3 (Short-chain dehydrogenase/reductase 3) is also named as retSDR1, DD83.1 and belongs to the short-chain dehydrogenases/reductases (SDR) family. It catalyzes the reduction of all-trans-retinal to all-trans-retinol in the presence of NADPH. It is detected high levels of expression in fetal kidney, liver, and lung and in adult heart, placenta, lung, liver, kidney, pancreas, thyroid, testis, stomach, trachea, spinal cord and lower levels in skeletal muscle, intestine, and lymph node. retSDR1 is barely detectable in adrenals, brain, thymus, and hematopoietic tissues (PMID:11861404). retSDR1 may play a more general role in retinol metabolism since its expression was observed in many fetal and adult tissues (PMID:20543567).

## Notable Publications

Author	Pubmed ID	Journal	Application
Lin Shen	36150660	Biochim Biophys Acta Mol Basis Dis	WB
Mark K. Adams	34542554	Biochem J	WB
Mark K Adams	27793605	Chem Biol Interact	WB

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 15393-1-AP (DHRS3 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 15393-1-AP (DHRS3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 15393-1-AP (DHRS3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using DHRS3 antibody (15393-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).



IP result of anti-DHRS3 (IP:15393-1-AP, 4ug; Detection:15393-1-AP 1:500) with A375 cells lysate 3600ug.