For Research Use Only

BDH1 Polyclonal antibody

Catalog Number:15417-1-AP

Featured Product





Basic Information	Catalog Number: GenBank / 15417-1-AP BC005844		Accession Number: Purification Method: Antigen affinity purification					
	Size:	GenelD (NCBI):	Recommended Dilutions: WB 1:2000-1:10000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate				
	500 ug/ml	622						
	Source: Rabbit	UNIPROT ID: Q02338						
	Isotype: IgG Immunogen Catalog Number: AG7705	Full Name: 3-hydroxybutyrate dehydrogenase, type 1 Calculated MW: 38 kDa		IHC 1:50-1:500 IF-P 1:50-1:500				
					Observed MW 31 kDa	t.		
					Applications	Tested Applications: WB, IHC, IF-P, IP, ELISA		
		Cited Applications: HT-29 cells,		cells, mouse liver tissue, rat liver tissue, human liver tissue, rat brain tissue, COL ouse brain tissue, mouse heart tissue				
·								
human, mouse, rat			IHC : rat brain tissue, human liver cancer tissue					
Cited Species:		IF-P: mouse	se liver tissue,					
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0								
Background Information	BDH1 (3-hydroxybutyrate dehydrogenase, type 1) is a 343 amino acid protein that localizes to the mitochondrial matrix and belongs to the short-chain dehydrogenases/reductases (SDR) family. The deduced 343 amino acid protein contains a 46 residue leader peptide, which is cleaved to produce the mature form. It is a lipid-requiring mitochondrial enzyme with a specific requirement of phosphatidylcholine (PC).							
Notable Publications	Author Pubmed ID Journal		Application					
	Nguyen Ngoc Trang	36184012	J Nutr Biochem	WB				
	De Huang	27644987	Cell Res	WB				
	Tim Düking	36112685	Sci Adv	WB				
Storage	Storage: Store at -20°C. Stable for one ye Storage Buffer: PBS with 0.02% sodium azide at Aliquoting is unnecessary for -2	nd 50% glycerol pH	7.3.					


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 15417-1-AP (BDH1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Various lysates were subjected to SDS PAGE followed by western blot with 15417-1-AP (BDH1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




mouse liver tissue were subjected to SDS PAGE followed by western blot with 15417-1-AP (BDH1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 15417-1-AP (BDH1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 15417-1-AP (BDH1 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-BDH1 (IP:15417-1-AP, 4ug; Detection:15417-1-AP 1:300) with mouse liver tissue lysate 4000ug.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse liver tissue using BDH1 antibody (15417-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).