For Research Use Only

MRPL49 Polyclonal antibody

Catalog Number: 15542-1-AP 6 Publications



Purification Method:

WB: 1:500-1:1000

Antigen affinity purification

IP: 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

Basic Information

Catalog Number:

15542-1-AP

BC004378

Concentration:

500 ug/ml

740

Source:

Rabbit

Q13405

Isotype:

GenBank Accession Number:

BC004378

GeneID (NCBI):

740

UNIPROT ID:

Q13405

Full Name:

Q13405 protein lysate
IHC: 1:50-1:500
Full Name:
mitochondrial ribosomal protein L49

IgG

AG7898

Immunogen Catalog Number: Calculated MW:

19 kDa

Observed MW: 19 kDa

Applications

Tested Applications: WB, IHC, IP, ELISA Cited Applications:

WB

Species Specificity:

human Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: BxPC-3 cells, A549 cells, HeLa cells

IP: HeLa cells,

IHC : human breast cancer tissue,

Background Information

MRPL49, also named as C11orf4, NOF1, or SW-cl.67, is a 166 amino acid protein, which belongs to the ribosomal protein L49em family. MRPL49 localizes in the Mitochondrion and is widely expressed in various tissues. MRPL49 is a regulator of C3 convertase of classical pathway.

Notable Publications

Author	Pubmed ID	Journal	Application
Joachim M Gerhold	26478270	Sci Rep	WB
Kazuo Yamamoto	24656129	Mol Cell	WB
Nina Rajala	24163258	Nucleic Acids Res	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 15542-1-AP (MRPL49 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 15542-1-AP (MRPL49 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



BxPC-3 cells were subjected to SDS PAGE followed by western blot with 15542-1-AP (MRPL49 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



IP result of anti-MRPL49 (IP:15542-1-AP, 4ug; Detection:15542-1-AP 1:500) with HeLa cells lysate 3360ug.