## For Research Use Only

# HSDL2 Polyclonal antibody

Catalog Number: 15631-1-AP

Featured Product

**6 Publications** 



**Basic Information** 

Catalog Number:

15631-1-AP

Size:

GeneID (NCBI):

600 µg/ml

84263

Source:

UNIPROT ID:

Rabbit

Q6YN16

Isotype:

GeneID (NCBI):

GeneID (NCBI):

GeneID (NCBI):

Full Name:

hydroxysteroid dehydrogenase like 2

Immunogen Catalog Number: Calculated MW:

AG7987 45 kDa

Observed MW: 45 kDa

**Applications** 

Tested Applications: IF/ICC, IHC, WB, ELISA Cited Applications: WB, IF, FC, IHC Species Specificity: human, mouse, rat

Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HEK-293 cells, HeLa cells

IHC: human liver cancer tissue, human prostate cancer

**Purification Method:** 

WB 1:1000-1:4000 IHC 1:50-1:500

IF 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

iccup

IF : HeLa cells,

# **Background Information**

Human hydroxysteroid dehydrogenase-like 2 (HSDL2) is a characterized SDR gene that not only catalyses the oxidation and reduction of multiple substrates but also regulates different metabolic and signalling pathways. Accumulating evidences suggest that HSDL2 play an important role in cancer progression. HSDL2 has promoting effects on tumor progression in papillary thyroid cancer, bladder cancer, ovarian cancer, and glioma, but has suppressing effects in cholangiocarcinoma . ( PMID: 31372054, PMID: 32211805)

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Bing Dong	31372054	Cancer Manag Res	IF
Songlin Zeng	25604190	Proteomics	WB
Anna Han	33488098	Onco Targets Ther	IHC,WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 15631-1-AP (HSDL2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 15631-1-AP (HSDL2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using HSDL2 antibody (15631-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).