For Research Use Only

OPA3 Polyclonal antibody

Catalog Number: 15638-1-AP

Featured Product

7 Publications



Basic Information

Catalog Number: GenBank Accession Number: 15638-1-AP BC005059
Size: GeneID (NCBI): 400 µ g/ml 80207
Source: UNIPROT ID: Rabbit Q9H6K4

Isotype: Full Name:
IgG optic atrophy 3 (autosomal recessive,

Immunogen Catalog Number: with chorea and spastic paraplegia)
AG8103 Calculated MW:

8103 Calculated MW: 179 aa, 20 kDa

> Observed MW: 20 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA Cited Applications:

WB, IHC, IP Species Specificity: human, mouse, rat Cited Species:

human, mouse, rat, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: A549 cells, human kidney tissue, mouse thymus tissue, Jurkat cells, HeLa cells, HEK-293T cells

Purification Method:

WB 1:500-1:2000

protein lysate

IHC 1:50-1:500

IF/ICC 1:50-1:500

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

IP: HeLa cells,

IHC: human liver tissue, mouse colon tissue

IF/ICC : HeLa cells,

Background Information

The OPA3 cDNA encodes a deduced 179-amino acid protein. Northern blot analysis demonstrated a primary transcript of approximately 5.0 kb that was ubiquitously expressed, most prominently in skeletal muscle and kidney. Mutations in this gene have been shown to result in 3-methylglutaconic aciduria type III and autosomal dominant optic atrophy and cataract.

Notable Publications

Author	Pubmed ID	Journal	Application
Miao Liu	35176941	Bioengineered	WB
Ning Meng	31881642	Cancers (Basel)	IHC
Alice E Navein	27106103	Hum Mol Genet	IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 15638-1-AP (OPA3 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human liver tissue slide using 15638-1-AP (OPA3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 15638-1-AP (OPA3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-OPA3 (IP:15638-1-AP, 4ug; Detection:15638-1-AP 1:1000) with HeLa cells lysate 1240 ug.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using OPA3 antibody (15638-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).