For Research Use Only

IKKB Polyclonal antibody

Catalog Number: 15649-1-AP

Featured Product

115 Publications

BC006231

3551

014920

GeneID (NCBI):

UNIPROT ID:

Full Name:



Basic Information

Catalog Number: 15649-1-AP Source: Rabbit

Isotype: IgG

Immunogen Catalog Number:

AG8191

inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase beta

Calculated MW: 756aa,81 kDa; 256aa,29 kDa

GenBank Accession Number:

Observed MW:

80 kDa, 86 kDa, 87 and 29 kDa

Applications

Tested Applications: WB, IHC, IP, ELISA Cited Applications: WB, IHC, IF, IP, CoIP Species Specificity: human, mouse, rat Cited Species:

human, mouse, rat, pig, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

WB: Jurkat cells, mouse brain tissue, rat brain tissue.

Purification Method:

WB: 1:1000-1:6000

protein lysate

IHC: 1:50-1:500

Antigen affinity purification

IP: 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

rat lung tissue IP: Jurkat cells.

Positive Controls:

IHC: human liver cancer tissue, human prostate cancer

Background Information

 $IKBKB, also \ named\ as\ IKKB, IKK2, NFKBIKB\ and\ IKK-B, belongs\ to\ the\ protein\ kinase\ superfamily, Ser/Thr\ protein\ and\ IKK-B,\ belongs\ to\ the\ protein\ kinase\ superfamily, Ser/Thr\ protein\ superfamily,$ kinase family and I-kappa-B kinase subfamily. IKBKB is a Serine kinase that plays an essential role in the NF-kappa-B signaling pathway. It acts as part of the canonical IKK complex in the conventional pathway of NF-kappa-B activation and phosphorylates inhibitors of NF-kappa-B on 2 critical serine residues. In addition to the NF-kappa-B inhibitors, IKBKB phosphorylates several other components of the signaling pathway including NEMO/IKBKG, NFkappa-B subunits RELA and NFKB1, as well as IKK-related kinases TBK1 and IKBKE. It also phosphorylates other substrates including NCOA3, BCL10 and IRS1. Within the nucleus, IKBKB acts as an adapter protein for NFKBIA degradation in UV-induced NF-kappa-B activation. This antibody can identify 4 isoform of IKBKB with the molecular weight of 80, 86, 87 and 29 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Xin Shen	36184549	Int Heart J	WB
Yanliang Wu	34601083	J Ethnopharmacol	WB
Wenbin Pei	34650433	Front Pharmacol	WB

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of IKKB antibody (15649-1-AP; 1:300; incubated at room temperature for 1.5 hours) with sh-Control and sh-IKKB transfected Jurkat cells.




Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 15649-1-AP (IKKB antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 15649-1-AP (IKKB antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 15649-1-AP (IKKB antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



IP result of anti-IKKB (IP:15649-1-AP, 4ug; Detection:15649-1-AP 1:3000) with Jurkat cells lysate 1400 ug.