

For Research Use Only

Hexokinase 1 Polyclonal antibody

Catalog Number: 15656-1-AP

9 Publications



Basic Information

Catalog Number:

15656-1-AP

Size:

240 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG8008

GenBank Accession Number:

BC008730

GeneID (NCBI):

3098

UNIPROT ID:

P19367

Full Name:

hexokinase 1

Calculated MW:

102 kDa

Observed MW:

100 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IHC 1:50-1:500

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IHC

Species Specificity:

human

Cited Species:

human, mouse, rat

Positive Controls:

WB : MCF-7 cells, U-87 MG cells

IHC : human stomach tissue,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

The hexokinases (HKs) catalyze the first obligatory step in glucose metabolism to generate glucose-6-phosphate. This not only furthers glucose entry by maintaining the concentration gradient for facilitated glucose influx but also provides the first intermediate for essentially all major pathways using glucose. There are four conventional HK isoforms, HK1/2/3/4, encoded by four different genes. Most adult tissues express only HK1. Muscle and adipose tissue use HK2 for glycolysis, liver, and pancreatic beta cells express HK4 (also called glucokinase) and do not express HK1 or HK2. (PMID: 31434645, PMID: 31848318)

Notable Publications

Author	Pubmed ID	Journal	Application
Hong Jin	33253911	Cell Signal	WB
Yanqing Li	33204111	Onco Targets Ther	WB
Li Yang	31332157	Med Sci Monit	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 15656-1-AP (Hexokinase 1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human stomach tissue slide using 15656-1-AP (Hexokinase 1 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).