For Research Use Only

CDC45L Polyclonal antibody

Catalog Number: 15678-1-AP

Featured Product

5 Publications



Basic Information

 Catalog Number:
 GenBank Accession Number:

 15678-1-AP
 BC010022

 Size:
 GeneID (NCBI):

 550 ug/ml
 8318

 Source:
 UNIPROT ID:

 Rabbit
 075419

Isotype: Full Name:
IgG CDC 45 cell division cycle 45-like (S.

Immunogen Catalog Number: Cerevisiae)
AG8150 Calculated MW:
64 kDa

Observed MW: 60-66 kDa Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:3000

IF/ICC 1:200-1:800

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB, IHC

Species Specificity: human, mouse Cited Species: human Positive Controls:

WB: K-562 cells, HEK-293 cells, HeLa cells, Jurkat cells,

mouse thymus tissue IP: K-562 cells,

IHC: mouse testis tissue,

IF/ICC: HeLa cells,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0.

buffer pH 6.0

Background Information

CDC45L(Cell division control protein 45 homolog), plays a role in the initiation of DNA replication, and genetically interacts with MCM family members and with ORC2 and physically form a complex including Mcm5p. It's the first gene missed in DiGeorge syndrome that suggest involve in cell proliferation. This is a rabbit polyclonal antibody raised against part of C-terminus of human CDC45L.

Notable Publications

Author	Pubmed ID	Journal	Application
Kimberlie A Wittig	34534348	Nucleic Acids Res	WB
Yue Wang	35414135	Signal Transduct Target Ther	WB
Lifei Fan	28467790	Oncotarget	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




K-562 cells were subjected to SDS PAGE followed by western blot with 15678-1-AP (CDC 45L antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




IP result of anti-CDC45L (IP:15678-1-AP, 3ug; Detection:15678-1-AP 1:1000) with K-562 cells lysate 3400ug.



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 15678-1-AP (CDC 45L antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 15678-1-AP (CDC 45L antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CDC 45L antibody (15678-1-AP) at dilution of 1:400 and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).