For Research Use Only

DLG5 Polyclonal antibody

Catalog Number:15687-1-AP 1 Publications




Basic Information	Catalog Number: 15687-1-AP	GenBank Accession Number: BC002915	Purification Method: Antigen affinity purification
	Size:	GenelD (NCBI):	Recommended Dilutions:
	400 ug/ml	9231	WB 1:500-1:1000
	Source: Rabbit	UNIPROT ID: Q8TDM6	IHC 1:50-1:500
	lsotype: IgG	Full Name: discs, large homolog 5 (Drosophila)	
	Immunogen Catalog Number: AG8194	Calculated MW: 1893 aa, 209 kDa	
		Observed MW: 75 kDa, 240 kDa	
Applications	Tested Applications:	Positive Controls:	
	WB, IHC, ELISA	WB: PC-3	cells, mouse kidney tissue, HeLa cells
	Cited Applications: WB	IHC : Human bowens disease,	
	Species Specificity: human, mouse		
	Cited Species: human		
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		
Background Information	Discs large homolog 5 (DLG5) belongs to the membrane-associated guanylate kinase (MAGUK) family. DLG5 participates in epithelial cell polarity maintenance and cancer development. DLG5 is highly expressed in normal tissues, but its expression is decreased in many cancers, such as bladder cancer, prostate cancer, breast cancer, and hepatocellular carcinoma (PMID: 32015336). Down-regulation of DLG5 is highly correlated with clinical tumor stage In breast cancer, knockdown of DLG5 induces cell migration, and overexpression of DLG5 inhibits cell migration (PMID: 28169360). DLG5 has 3 isoforms with the molecular weight of 75, 214 and 240 kDa.		
Notable Publications	Author	Pubmed ID Journal	Application
	Rui Su	32015336 Cell Death Dis	WB
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	50% glycerol pH 7.3.	

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data

1.5 hours.




PC-3 cells were subjected to SDS PAGE followed by western blot with 15687-1-AP (DLG5 antibody) at dilution of 1:500 incubated at room temperature for



Various lysates were subjected to SDS PAGE followed by western blot with 15687-1-AP (DLG5 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human brown disease slide using 15687-1-AP (DLG5 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human brown disease slide using 15687-1-AP (DLG5 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).