

For Research Use Only

SMPX Polyclonal antibody

Catalog Number: 15791-1-AP



Basic Information

Catalog Number: 15791-1-AP	GenBank Accession Number: BC005948	Purification Method: Antigen affinity purification
Size: 500 µg/ml	GeneID (NCBI): 23676	Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500
Source: Rabbit	UNIPROT ID: Q9UHP9	
Isotype: IgG	Full Name: small muscle protein, X-linked	
Immunogen Catalog Number: AG8497	Calculated MW: 88 aa, 10 kDa	
	Observed MW: 9 kDa	

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

human, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : pig skeletal muscle tissue, human skeletal muscle tissue, human colon tissue, pig colon tissue

IHC : human kidney tissue, human brain tissue, human lung tissue, human ovary tissue, human spleen tissue, human testis tissue

Background Information

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:

T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 15791-1-AP (SMPX antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 15791-1-AP (SMPX antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 15791-1-AP (SMPX antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



pig skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 15791-1-AP (SMPX antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.