For Research Use Only

GSTP1 Polyclonal antibody

Catalog Number: 15902-1-AP

Featured Product

38 Publications



Basic Information

Catalog Number: 15902-1-AP

GenBank Accession Number: BC010915

Purification Method: Antigen affinity purification

Size:

GeneID (NCBI): 2950

Recommended Dilutions: WB 1:2000-1:16000

600 ug/ml Source: Rabbit

UNIPROT ID: P09211

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Isotype:

Full Name: glutathione S-transferase pi 1 protein lysate IHC 1:200-1:1600

IF/ICC 1:50-1:500

Immunogen Catalog Number:

Calculated MW: AG8731 210 aa, 23 kDa

> Observed MW: 23-28 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), IP, ELISA

Cited Applications: WB, IHC, IF, IP, CoIP

Species Specificity: human, mouse, rat **Cited Species:**

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: COLO 320 cells, human brain tissue, HeLa cells, HEK-293 cells, K-562 cells, Jurkat cells, PC-3 cells, mouse brain tissue, mouse heart tissue, rat brain tissue, rat heart tissue

IP: mouse brain tissue,

IHC: mouse skin tissue, human colon, human liver

tissue, human skin tissue

IF/ICC: HepG2 cells,

Background Information

Glutathione S-transferase P (GSTP1) conjugates reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles. It is involved in the formation of glutathione conjugates of both prostaglandin A2 (PGA2) and prostaglandin J2 (PGJ2) (PubMed:9084911). Participates in the formation of novel hepoxilin regioisomers (PubMed:21046276).

Notable Publications

Author	Pubmed ID	Journal	Application
Lei Ye	33491741	Int J Oncol	WB
Shuo Chen	26396496	Drug Des Devel Ther	WB
Ming-Wei Lin	36135183	Curr Issues Mol Biol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 15902-1-AP (GSTP1 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




WB result of GSTP1 antibody (15902-1-AP; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-GSTP1 transfected HeLa cells.



IP result of anti-GSTP1 (IP:15902-1-AP, 3ug; Detection:15902-1-AP 1:800) with mouse brain tissue lysate 4000ug.



Immunohistochemical analysis of paraffinembedded mouse skin tissue slide using 15902-1-AP (GSTP1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




1x10^6 HepG2 cells were intracellularly stained with 0.4 ug GSTP1 Polyclonal antibody (15902-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L) (SA00013-2)(red), or 0.4 ug Rabbit 1gG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunohistochemical analysis of paraffinembedded mouse skin tissue slide using 15902-1-AP (GSTP1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon slide using 15902-1-AP (GSTP1 antibody) at dilution of 1:1600 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using GSTP1 antibody (15902-1-AP) at dilution of 1:200 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).