

For Research Use Only

IDH3A Polyclonal antibody

Catalog Number: 15909-1-AP

Featured Product

17 Publications



Basic Information

Catalog Number:

15909-1-AP

Size:

600 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG8662

GenBank Accession Number:

BC021967

GeneID (NCBI):

3419

UNIPROT ID:

P50213

Full Name:

isocitrate dehydrogenase 3 (NAD+)

alpha

Calculated MW:

366 aa, 40 kDa

Observed MW:

40 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:5000-1:50000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:20-1:200

IF 1:20-1:200

Applications

Tested Applications:

FC, IF/ICC, IHC, IP, WB, ELISA

Cited Applications:

IF, IHC, IP, WB

Species Specificity:

human, mouse, rat

Cited Species:

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HepG2 cells, human skeletal muscle tissue, mouse brain tissue, rat brain tissue

IP: HepG2 cells,

IHC: human brain tissue,

IF: HepG2 cells,

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Yingjiang Xu	34704268	EMBO J	WB
Teiji Oda	26437904	Circ J	WB
Péter Gulyássi	33211194	Amino Acids	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 15909-1-AP (IDH3A antibody) at dilution of 1:1200 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human brain using 15909-1-AP (IDH3A antibody) at dilution of 1:100 (under 40x lens).




Immunohistochemical analysis of paraffin-embedded human brain using 15909-1-AP (IDH3A antibody) at dilution of 1:100 (under 10x lens).




IP result of anti-IDH3A (IP:15909-1-AP, 4 μ g; Detection:15909-1-AP 1:1000) with HepG2 cells lysate 2400 μ g.




Immunofluorescent analysis of HepG2 cells using 15909-1-AP (IDH3A antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 15909-1-AP (IDH3A antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using IDH3A antibody (15909-1-AP) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).



1X10⁶ HepG2 cells were intracellularly stained with 0.4 μ g Anti-Human IDH3A (15909-1-AP) and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 μ g Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).