For Research Use Only

## TNFAIP8L2 Polyclonal antibody

Catalog Number:15940-1-AP

Featured Product

37 Publications




Basic Information	Catalog Number: 15940-1-AP	GenBank Accession Number: BC063014		Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):		Recommended Dilutions:	
	400 ug/ml	79626		WB 1:1000-1:6000	
	Source:	UNIPROT ID:		IP 0.5-4.0 ug for 1.0-3.0 mg of total	
	Rabbit	Q6P589 Full Name:		protein lysate IHC 1:50-1:500 IF/ICC 1:50-1:500	
	Isotype:				
	IgG	tumor necrosis factor, alpha-induced protein 8-like 2 Calculated MW: 184 aa, 21 kDa Observed MW: 18-20 kDa			
	Immunogen Catalog Number: AG7964				
	Applications				Tested Applications:
WB, IHC, IF/ICC, FC (Intra), IP, ELISA			WB: THP-1 ce	WB : THP-1 cells, mouse small intestine tissue, mouse	
Cited Applications: cerebel		cerebellum ti	m tissue, rat lymph tissue, rat spleen tissue 7 cells, mouse spleen tissue		
Species Specificity:			IP : mouse spl	een tissue,	
human, mouse, rat			IHC : human lymphoma tissue, human ovary tumor		
Cited Species: tissue, mou human, mouse, rat		tissue, mouse	se spleen tissue		
Note-IHC: suggested antigen retrieval with <b>TE buffer pH 9.0; (*)</b> Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0					
Background Information	TIPE2 (TNF- α -inducible protein 8-like 2, or TNFAIP8L2), a negative immune regulator, is essential for maintaining immune homeostasis. TIPE2 shares considerable sequence homology with members of the tumor necrosis factor- α inducible protein 8 (TNFAIP8) family, which are thought to regulate cellular and immune homeostasis. TIPE2 acts a a negative regulator of Toll-like receptor and T-cell receptor function and is preferentially expressed in lymphoid tissues. TIPE2 plays an essential role for a signal transduction pathway that links inflammatory immune response to specific conditions after cerebral ischemia				
Notable Publications	Author Pu	ıbmed ID Jo	ournal	Application	
	Jiacheng Bi 34	4524845 S	ci Adv	WB	
		7601471 J	Biol Chem	WB,IF	
	Svetlana A Fayngerts 29	9058702 N	lat Immunol	WB,IF,IP	
Storage	Storage: Store at -20°C. Stable for one year at	ter shipment.			

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data


temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 15940-1-AP (TIPE2 antibody) at dilution of 1:3000 incubated at room




Immunohistochemical analysis of paraffinembedded human lymphoma tissue slide using 15940-1-AP (TIPE2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human lymphoma tissue slide using 15940-1-AP (TIPE2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-TIPE2 (IP:15940-1-AP, 4ug; Detection:15940-1-AP 1:2000) with mouse spleen tissue lysate 1400 ug.



Immunofluorescent analysis of (4% PFA) fixed RAW 264.7 cells using TIPE2 antibody (15940-1-AP) at dilution of 1:200 and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10^6 THP-1 cells were intracellularly stained with 0.4 ug Anti-Human TIPE2 (15940-1-AP) and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).