## For Research Use Only

# **BAAT Polyclonal antibody**

Catalog Number: 15990-1-AP

1 Publications



**Purification Method:** 

WB 1:500-1:1000 IHC 1:20-1:200

IF/ICC 1:10-1:100

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

Catalog Number: 15990-1-AP Size:

500 µ g/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG8860

GenBank Accession Number: BC009567

GeneID (NCBI): 570 UNIPROT ID:

Q14032 Full Name:

bile acid Coenzyme A: amino acid Nacyltransferase (glycine Ncholoyltransferase)

Calculated MW: 418 aa, 46 kDa Observed MW: 50 kDa

**Applications** 

Tested Applications: IF/ICC, IHC, WB, ELISA Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

human, mouse

Positive Controls:
WB: human liver tissue,

IHC: human hepatocirrhosis tissue,

IF/ICC: HepG2 cells,

# **Background Information**

BAAT is bile acid-CoA:amino acid N-acyltransferase which involved in in bile acid metabolism. BAAT acts as an acyl-CoA thioesterase that regulates intracellular levels of free fatty acids.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Chatterjee Sagnik S	24211540	Toxicol In Vitro	

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


### **Selected Validation Data**




human liver tissue were subjected to SDS PAGE followed by western blot with 15990-1-AP (BAAT antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human hepatocirrhosis tissue slide using 15990-1-AP (BAAT Antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human hepatocirrhosis tissue slide using 15990-1-AP (BAAT Antibody) at dilution of 1:100 (under 40x lens).



Immunofluorescent analysis of HepG2 cells using 15990-1-AP (BAAT antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).