## For Research Use Only

# MMS19 Polyclonal antibody

Catalog Number: 16015-1-AP

Featured Product

17 Publications



**Basic Information** 

**Applications** 

Catalog Number: 16015-1-AP Size: 500 μg/ml

Rabbit Isotype:

Source:

Immunogen Catalog Number:

AG8792

**Tested Applications:** 

IF/ICC, WB, ELISA Cited Applications:

WB, CoIP

Species Specificity: human, mouse, rat Cited Species: human, mouse

GenBank Accession Number:

BC006575 GeneID (NCBI): 64210 **UNIPROT ID:** Q96T76 Full Name:

MMS19 nucleotide excision repair

homolog (S. cerevisiae)

Calculated MW: 1030 aa. 113 kDa Observed MW:

103-113 kDa

Positive Controls:

WB: HEK-293 cells, human brain tissue, PC-3 cells

**Purification Method:** 

WB 1:1000-1:6000 IF/ICC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

IF/ICC: HeLa cells,

# **Background Information**

MMS19 plays a critical role in the biogenesis of multiple Fe-S proteins, which functions in genomic stability, RNA  $polymerase\ II\ function, and\ telomere\ length\ regulation.\ MMS19\ has\ a\ alanine-\ and\ leucine-rich\ N\ terminus.\ It$ possible has a role in nucleotide excision repair(NER) and RNA polymerase II(POL II) transcription by interacting with ERCC2/XPB helicases, both subunits of NER-transcription factor TFIIH. Also MMS19 involves in the regulation of ER. It can function in chromosome segregation by acting as a part of the mitotic spindle-associated MMXD complex.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Seung Wook Yang	33004795	Nat Commun	WB
Xiaorui Fan	35654137	J Biol Chem	WB
Laura Mariotti	32576938	Commun Biol	WB


Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 16015-1-AP (MMS19 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



human brain tissue were subjected to SDS PAGE followed by western blot with 16015-1-AP (MMS19 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using 16015-1-AP (MMS19 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).